

Publication number:

**0 257 993**. A2

æ

## **EUROPEAN PATENT APPLICATION**

2 Application number: 87307384.5

Date of filing: 20.08.87

(5) Int. Cl.4: C 07 H 21/04 C 12 N 15/00, C 12 N 9/88, C 12 N 1/20, A 01 H 1/00, A 01 H 5/10, A 01 N 25/00, A 01 N 47/36

30 Priority: 26.08.86 US 900609

43 Date of publication of application: 02.03.88 Bulletin 88/09

Designated Contracting States:
 AT BE CH DE ES FR GB GR IT LI LU NL SE

(7) Applicant: E.I. DU PONT DE NEMOURS AND COMPANY 1007 Market Street Wilmington Delaware 19898 (US)

(2) Inventor: Bedbrook, John Robert 110 Cambridge Way Pledmont, California 94611 (US) Chaleff, Roy Scott 1 Spring Lane . Chadds Ford, Pennsylvania 19317 (US)


Falco, Saverio Carl 1902 Millers road Arden, Delaware 19810 (US)

Mazur, Berbara Jean 4601 Weldin road Wilmington, Delaware 19803 (US)

Yadav, Narendra Singh 127 Jade Drive Wilmington, Delaware 19810 (US)

Representative: Hildyard, Edward Martin et al Frank B. Dehn & Co. Imperial House 15-19 Kingsway London WC2B 6UZ (GB)

- (A) Nucleic acid fragment encoding herbicide resistant plant acetolactate synthese.
- A nucleic acid fragment encoding a herbicide-resistant plant acetolactate synthase protein is disclosed. This nucleic acid fragment contains at least one nucleotide mutation resulting in one amino acid change in one of seven substantially conserved regions of acetolactate synthase amino acid homology. This mutation results in the production of an acetolactate synthase protein which is resistant to sulfornylurea herbicide compounds compared to the wild-type protein. Transformation of herbicide sensitive plants or plant cells with the fragment results in resistance to the herbicide.



Bundesdruckerei Berlin

#### Description

## NUCLEIC ACID FRAGMENT ENCODING HERBICIDE RESISTANT PLANT ACETOLACTATE SYNTHASE

The present invention relates to nucleic acid fragments encoding a herbicide-resistant form of the enzyme acetolactate synthase (ALS).

Sulfonylurea herbicides such as sulforneturon methyl (I) and chlorsulfuron (II) inhibit growth of some bacteria, yeast and higher plants by blocking acetolactate synthase [ALS, EC 4.1.3.18], the first common enzyme in the biosynthesis of the branched-chain amino acids valine, leucine and isoleucine. The biosynthesis of branched-chain herbicides is restricted to plants and microbes. ALS is also inhibited by a structurally unrelated class of herbicides, the imidazolinones.

10

20

35

40

Three major isozymes of ALS, designated I, II and III, have been identified in enteric bacteria. Isozymes I and III, but not II, are sensitive to end-product inhibition by valine. Each of the three bacterial isozymes comprises a large and a small protein subunit. ALS enzymes from the yeast Saccharomyces cerevisiae and from some higher plants have been partially characterized and show some degree of end-product inhibition. It is not known if the yeast and plant ALS enzymes consist of one or more different polypeptides of the yeast and plant ALS enzymes are in the mitochondria and chloroplasts, respectively.

60

Genes encoding ALS enzymes have been isolated from the enteric bacteria Salmonella typhinurium and Escherichia Coli, and the yeast S. cerevisiae. The nucleotide sequences of the genes coding for the two subunits of E. coli ALS isozymes I, II and III show that they are organized as operons itvBN, ilvGM and iivIH, respectively. Comparison of the deduced amino acid sequences of the large subunits of the E. coli ALS isozymes shows three regions with about 50% conserved amino acids, comprising about two-thirds of the proteins, and separated by regions sharing little discernible homology. Amino acid sequence conservation, though less extensive, is also evident among the small subunits of the bacterial isozymes. In the yeast S. cerevisiae, a single gene, ILV2, essential for ALS activity was identified. Nucleotide sequence analysis of the ILV2 gene was revealed that the polypeptide encoded by it is homologous to the large subunits of the bacterial ALS isozymes. The deduced amino acid sequence of the yeast ALS shows the same degree of structural organization and the same degree of homology as it observed between the large subunits of the bacterial isozymes, except for about ninety amino acids at the amino terminus of the yeast protein that are believed to be involved in the translocation of the protein into the mitochondrion. No information on the structure of the plant genes encoding ALS or the amino acid sequence of plant ALS enzymes was available prior to the inventions disclosed herein.

Enteric bacterial isozyme I is the only ALS in nature that Is known to be insensitive to inhibition by sulformeturon methyl and chlorsulfuron. Therefore, enteric bacteria are sensitive to these herbicides only in the presence of valine, which inhibits isozyme I. Sulfonylurea herbicide-resistant mutant forms of the enteric bacteria Salmonella typhimurium and E. coli (selected in the presence of valine), the yeast S. cerevisiae and the higher plants Nicotaiana tabacum (tobacco), Arabidopsis thaliana and Zea mays (corn) have been identified. These mutant phenotypes cosegregate with herbicide-resistance forms of ALS through genetic crosses. In S.

typhimurium the herbicide-resistance mutations are genetically linked to a gene encoding ALS, and in <u>E. coli</u> and <u>S. cerevisiae</u>, these mutations reside in the structural genes for ALS. In the higher plants the mutations responsible for the resistance are inherited as single, dominant or semidominant nuclear traits. In tobacco, these mutations map to either of two unlinked genetic loci.

The chemical control of undesirable weeds associated with agronomically useful crops requires the use of highly selective chemical herbicides. In some cases, it is difficult to identify any chemical which kills weeds without injury to the crop plant. The introduction of herbicide-resistance as a biological trait in crop plants would overcome this difficulty.

5

15

35

45

*5*5

60

65

Although many genes involved in the structure and function of differentiated plant tissues and organs are not expressed in undifferentiated tissues, those involved in basic cellular functions are expressed and can be selected for in a disorganized callus or cell suspension culture. This has been demonstrated in many cases by the selection of a phenotype in tissue culture from which plants expressing the same phenotype have been regenerated. Examples include the in vitro selection of plants resistant to herbicides, pathotoxins or diseases, antibiotics, amino acid analogues, salt tolerance, etc. (reviewed in Chaleff, 1981; Evans et al, 1983; Vasil, 1986).

Since acetolactate synthase is an enzyme involved in the basic cellular metabolic activity of amino acid biosynthesis, it was expected and has been demonstrated that genes encoding this enzyme are expressed in callus tissue as well as the whole plant. The sulfonylurea resistant tobacco mutants described in this patent, S4, C3 and Hra, were first selected in tissue culture and subsequently regenerated into whole plants in which the resistant phenotypes were retained in a genetically stable manner (Chaleff and Ray, 1984). Callus tissues derived from regenerated plants or their progeny continue to grow on concentrations of the herbicide which inhibit the growth of wild type callus. Thus resistance to a sulfonylurea herbicide at the plant cellular level is predictive of resistance at the whole plant level. In addition, it has been demonstrated in bacteria, yeast and higher plants that mutations resulting in the production of herbicide resistant ALS are sufficient to confer resistance at the cellular level and, in the case of plants, at the whole plant level. Therefore, the observation of herbicide-resistant ALS in extracts of plant cells is also predictive of herbicide resistant growth of culture plant cells and herbicide resistant growth of whole plants.

There are limitations to obtaining herbicide-resistant crop plants through tissue culture or seed mutagenesis: 1) the tissue culture method is restricted to those crop species that can be manipulated in and regenerated from tissue culture, 2) plants derived from seed mutagenesis, and possibly from tissue culture, might have non-desirable phenotypes which would require multiple genetic backcrosses to eliminate, and 3) transfer of the resistance gene by breeding would be restricted to cultivars of the same species as well as require multiple genetic back crosses. Therefore, the isolation of a nucleic acid fragment able to confer herbicide resistance and its subsequent introduction into crops through genetic transformation should allow rapid, cross-species transfer of herbicide resistance and overcome many of the above mentioned limitations.

Although genes isolated from one plant have been introduced and expressed in other plants, non-plant genes have been expressed in plants only as chimeric genes in which the encoding sequences of the non-plant genes have been fused to plant regulatory sequences required for gene expression. However, it would be difficult to introduce herbicide resistance into plants by introducing chimeric genes consisting of bacterial or yeast genes for herbicide-resistant forms of ALS, since (a) these microbial ALS enzymes are believed to lack a specific signal (transit) peptide sequence required for uptake into plant chloroplasts, the cellular location of plant ALS, (b) the bacterial isozymes consist of two different polypeptide subunits, and (c) the microbial ALS enzymes may not function optimally in the foreign cellular environment of higher plants. Therefore, there is a need for nucleic acid fragments (1) which encode a herbicide-resistant form of plant ALS, and (2) which can confer herbicide resistance when introduced into herbicide sensitive plants.

According to one aspect of the present invention, there is provided a nucleic acid fragment comprising a nucleotide sequence encoding plant acetolactate synthase with increased resistance to one or more suffonylurea herbicides. The nucleotide sequence comprises at least one sequence which encodes one of the substantially conserved amino acid sub-sequences designated A, B, C, D, E, F and G in Figure 6. The nucleic acid fragment is further characterized in that at least one of the following conditions is met;

- a) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence A wherein  $\epsilon_1$  is an amino acid other than alanine, and/or  $\epsilon_2$  is an amino acid other than glycine,
- b) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence B wherein  $\alpha_1$  is an amino acid other than proline,
- c) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence C wherein  $\delta_2$  is an amino acid other than alanine,
- d) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence D wherein  $\lambda_1$  is an amino acid other than lysine.
- e) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence E wherein  $\gamma_1$  is an amino acid other than aspartic acid,
- f) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence F wherein  $\beta_3$  is an amino acid other than the tryptophan, and/or  $\beta_8$  is an amino acid other than valine and/or  $\beta_7$  is an amino acid other than phenylalanine, and
- g) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence G wherein  $\sigma_1$  is an amino acid other than methionine.

In another embodiment, the instant invention provides a nucleic acid fragment encoding plant acetolactate

synthase which is capable of being incorporated into a nucleic acid construct suitable for transforming a plant containing wild-type acetolacetate synthase which is sensitive to a sulfonylurea herbicide compound, said nucleic acid fragment having at least one point mutation relative to the wild-type nucleic acid fragment encoding plant acetolactate synthase such that upon transformation with said nucleic acid construction said plant contains and nucleic acid fragment and exhibits increased resistance to the application of said sulfonylurea herbicide compound.

In another embodiment, the present invention provides an acetolactate synthase protein which is resistance to a sulfonylurea herbicide compound comprising an amino acid sequence wherein a substitution of at least one amino acid has occurred.

In still another embodiment, the present invention provides nucleic acid constructs, monocotyledonous and dicotyledonous plants, and tissue cultures which contain the specified nucleic acid fragment. The invention further provides methods for tranforming plants with the specified fragments, selecting transformed plant cells, and growing transformed plants.

The present invention also provides a method for selecting plant cells transformed with the nucleic acid fragment of the present invention. The method comprises introducing the fragment into plant cells whose growth is sensitive to inhibition by herbicides to which the ALS encoded by the fragment is resistant to form a transformed plant cell. The transformed plant cells whose growth is resistant to the selected herbicide are identified by selection at a herbicide concentration which inhibits the growth of the untransformed plant cells.

In another aspect, the present invention is a method for controlling unwanted vegetation growing at a locus where a herbicide-resistant, agronomically useful plant (transformed with the nucleic acid fragment of the present invention) has been cultivated. The method comprises applying to the locus to be protected an effective amount of herbicide. In still another aspect, the present invention provides a nucleic aspect, the present invention provides a nucleic acid construct wherein a nucleic acid fragment encoding plant acetolactate synthase with increased sulfonylurea herbicide resistant as hereinbefore described is linked to a second nucleic acid fragment conferring a second trait such that when said nucleic acid construct is utilized to transform a plant, the expression of herbicide resistance by said plant upon application of sulfonylurea compounds can be utilized to detect the presence of said second nucleic acid fragment in said plant.

#### Brief Description of the Drawings

25

30

35

40

Figure 1 is a physical map of nucleic acid insert fragments containing ALS genes isolated from a genomic library of DNA from the tobacco Hra mutant.

Figure 2 is a diagram of plasmid pAGS152 showing a physical map of the nucleic acid fragment from tobacco encoding a herbicide-resistant ALS.

Figure 3 is a physical map of a nucleic acid insert fragment in phage clone 35 isolated from genomic library of DNA from the tobacco C3 mutant.

Figure 4 is a nucleotide sequence, and the cognate deduced amino acid sequence, of a gene from the Hra mutant of tobacco encoding a herbicide-resistant form of ALS from tobacco.

Figure 5 is a nucleotide sequence, and the cognate deduced amino acid sequence, of a gene from the C3 mutant of tobacco encoding a herbicide-resistant form of ALS.

Figure 6 is a comparison of deduced amino acid sequence of the large subunits of bacterial ALS and the yeast and plant ALS enzymes.

Figure 7 is a physical map of a nucleic acid insert fragment and sub-fragments derived from phage clone 21 isolated from genomic library of sugarbeet DNA.

Figure 8 is a comparison of deduced amino acid sequences of plant ALS enzymes.

The present invention provides specified nucleic acid fragments which confer herbicide resistance when Introduced into herbicide-sensitive plants. As used herein, the term "nucleic acid fragment" refers to a linear segment of single- or double-stranded deoxyribonucleic acid (DNA) or ribonucleic acid (RNA), which can be derived from any source. Preferably, the nucleic acid fragment of the present invention is a segment of DNA. The term "plant" refers to a photosynthetic organism including algae, mosses, ferns, gymnosperms, and angiosperms. The term, however, excludes, prokaryotic and eukaryotic microorganisms such as bacteria, yeast, and fungi. "Plant cell" includes any cell derived from a plant, including undifferentiated tissue such as callus or gall tumor, as well as protoplasts, and embryonic and gametic cells. The term "plant acetolactate synthase refers to the specified enzyme when expressed in a plant or plant cell. The term "nucleotide sequence" refers to a polymer of DNA or RNA which can be single- or double-stranded, optionally containing synthetic, non-natural, or altered nucleotides capable of incorporation into DNA or RNA polymers. As used herein, the expression "substantially conserved amino acid sequences" refers to regions of amino acid homology between polypeptides comprising ALS enzymes from different sources. In the present invention seven substantially conserved amino acid sequences, designated A, B, C, D, E, F, and G are shown in Figure 6. One skilled in the art could align the amino acid sequnces of ALS enzymes from different sources to the schematic of Figure 6 to identify the segments therein which are the substantially conserved amino acid sequences defined herein. The skilled person could then determine whether the identified segments have the characteristics disclosed and claimed in the present application. It is to be understood that the expression includes modification of the segments which do not adversely affect the activity of the ALS enzyme. The term "nucleic acid construct" refers to a plasmid, virus, autonomously replicating sequence, phage or linear segment of a single- or double-stranded DNA or RNA, derived from any source, which is capable of

A

introducing a nucleic acid fragment into a biological cell.

\*Regulatory nucleotide sequence\*, as used herein, refers to a nucleotide sequence located 5' and/or 3' to a nucleotide sequence whose transcription and expression is controlled by the regulatory nucleotide sequence in conjuction with the protein synthetic apparatus of the cell. As used herein, a \*regulatory nucleotide sequence\* can include a promoter region, as that term is conventionally employed by those skilled in the art. A promoter region can include an association region recognized by an RNA polymerase, one or more regions which control the effectiveness of transcription initiation in responst to physiological consitions, and a transcription initiation sequence.

"Transit peptide" refers to a signal polypeptide which is translated in conjunction with a polypeptide which by a product nucleotide sequence, forming a polypeptide precursor. In the process of transport to a selected site within the cell, for example, a chloroplast, the transit peptide can be cleaved from the remainder of the polypeptide precursor to provide an active or mature protein.

10

15

35

55

60

65

Herbicide," as used herein, refers to an antiblotic compound which inhibits the metabolism, growth, or replication of plant cells or whole plants. Cells transformed with a construct of the present invention exhibit selectable cross-resistance to certain structurally related sulfonamide compounds effective as broad-spectrum preemergent and postemergent herbicides. As used herein in a generic sense, "sulfonylurea herbicides" refer to N-(heterocyclicaminocarbonyl)arylsulfonamide compounds exhibiting broad-spectrum herbicidal activity and low mammalian toxicity "Selective concentration" refers to a concentration of an inhibitor or antibiotic compound, for example, a herbicide, which is capable of inhibiting the metabolism, growth, or multiplication of wild-type cell or organism. Such an organism, as well as clones thereof, is referred to as a "sensitive" organism or cell. "Resistance" refers to a capability of an organism or cell to grow in the presence of selective concentrations of an inhibitor. In relation to particular enzymes or proteins, "sensitive" indicates that the enzyme or protein is susceptible to specific inhibition by a particular inhibiting compound, for example, an antibiotic or herbicide. In relation to particular enzymes or proteins, "resistant" indicates that the enzyme or protein, as a result of a different chemical structure, expresses activity in the presence of a selective concentration of a specific inhibitor which inactivates sensitive variants of the enzyme or protein. The term "selectable genetic market" refers to a nucleotide sequence which, when incorporated into the genome of an organism, allows growth of that organism and its progeny under conditions which inhibit growth of the organism lacking the selectable genetic marker. For example, a gene which encodes an enzyme that is resistant to specific inhibition by a particular antibiotic compound, such as a herbicide, can function as a selectable genetic marker by allowing an organism, such a plant, to grow and propagate in the presence of a selective concentration of the compound. A second nucleic acid fragment, controlling a property which is difficult to assay, can be covalently linked to the selectable genetic marker, in which case the presence of the selectable marker, indicated by growth of an organism under selective conditions, can be used to detect an organism containing the second nucleic acid fragment.

Preparation of DNA Fragments Encoding Herbicide-Resistant ALS

Callus cultures of sensitive tobacco (Nicotiana tabacum var. Xanthi) were exposed to sulfometuron methyl at 2 ppb according to the method described by Chaleff, U.S. 4,443,971. Resistant cell lines designated C3 and S4 were selected. Standard genetic analysis of plants regenerated from these cells lines indicated that the C3 and S4 lines each carried a single semi-dominant nuclear gene mutation responsible for the herbicide resistance trait and that the C3 and S4 mutations were not genetically linked, i.e. were in different genes designated SURA and SURB, respectively. The C3 and S4 lines were shown to produce ALS enzyme activity one-hundred fold more resistance to the sulfonylurea herbicides chlorsulfuron and sulfomenturon methyl than ALS from wild type. Production of herbicide resistance ALS activity cosegregated in genetic crosses with resistance to growth inhibition by the herbicides. The observation of two different genes that had mutated to form herbicide resistance ALS was not unexpected because N. tabacum is believed to be an allotetraploid plant formed from N. tomentosiformis and N. sylvestris, essentially containing two complete genomes. Thus, the S4 and C3 cell lines each contain one mutant and one wild type ALS gene. The S4 cell line was exposed to sulforneturon methyl at 200 ppb, a selective concentration which completely inhibits the growth of S4. Cell lines resistant to 200 ppb were identified; one such line was designated Hra. Hra was shown to tolerate concentrations of sulfometuron methyl one thousand times greater than that required to completely inhibit the growth of wild type callus. Hra was shown to be cross resistant to chlorsulfuron. Plants were regenerated from Hra callus cultures. Genetic analysis of the plants demonstrated that the Hra and S4 mutations were linked indicating that the Hra line contained a second mutation in the mutant gene of the progenitor S4 line. ALS activity in extracts of leaves of wild type and homozygous Hra mutant tobacco plants was determined. The ALS activity in the extract from Hra mutant plants was about one thousand fold more resistant to chlorsulfuron than was the activity of the wild type plants. Hra mutant plants were further shown to be cross resistant to the foliar application of the following compounds:

- 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-2-pyridinecarboxylic acid. (1-methylethanamine) salt:
- 5-ethyl-4,5-dihydro-2-[4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid;
- 2-(2-chloroethoxy)-N-[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)aminocarbonyl]benzenesulfonamide;
- 2-chloro-N-[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)aminocarbonyl]benzenesulfonamide;
- 2-[[(4-chloro-6-methoxypyrimidin-2-yl)aminocarbonyl]aminosulfonyl]benzoic acid, ethyl ester;

- N-[(4,6-dimethoxypyrimidin-2-yl)aminocarbonyl]-2,3-dihydro-2-methylbanzo[β]thiophene-7-sulfonamide,
   1.1-dioxida:
- 7-chloro-N-[(4,6-dimethoxypyrimidin-2-yl)aminocarbonyl]-3,4-dihydro-2-methyl-2H-1,2-benzothlazine-8-sul-fonamide, S,5-dioxide;
- 2-[[(4-methoxy-6-methylpyrimidin-2-yl)aminocarbonyl]aminosulfonyl]-6-methylbenzolc acid, methyl ester;
- 5,7-dimethyl-N-(2-methyl-6-nitrophenyl)[1,2,4]triazolo[1,5-A]pyrimidin-2-sulfonamide;
- 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid;
- 6-(4-isopropyl-4-methyl-5-oxo-2-imidazolin-2-yl)-m-toluic acid and p-toluic acid, methyl esters;
- 2-[[(4,6-dimethylpyrlmldln-2-yl)aminocarbonyl]aminosulfonyl]benzoic acid, methyl ester;
- N-(2,6-dichlorophenyl)-5,7-dimethyl[1,2,4]trlazolo [1,5-A]pyrimidin-2-sulfonamide;

30

35

45

• N-(2-chloro-6-methylphenyl)-5,7-dimethyl[1,2,4]triazolo[1,5-A]pyrimidin-2-sulfonamide.

In order to clone a herbicide resistant ALS gene, tobacco DNA was isolated from the S4 homozygous mutant line of Nicotiana tabacum. 50 g portions of callus tissue were frozen in liquid N2, and then lyophilized. The resulting dried tissue was then ground at about 23°C in a blender, using 15 second bursts, until powdered. Ten volumes of a sucrose buffer (0.3 M sucrose, 50 mM Tris-HCl pH 8.0, 5mM MgCl2) were added, and the resulting suspension was incubated at 0°C for 5 minutes. The suspension was then filtered through cheesecloth, and centrifuged at 350 x g for 10 minutes. The nuclear pellet was then resuspended in lysis buffer (20 mM EDTA, 50 mM Tris-HCl pH 8.0, 1% Sarkosyl), CsCl added to provide 0.95 g per mL. buffer, and the resulting mixture centrifuged at 17,000 x g for 20 minutes at 4°. Ethicium bromide was added to the resulting supernatant to a concentration of 400 µg per mL, the reflective index was adjusted to 1.39, and the resulting solution centrifuged at 90,000 x g in a Beckman Ti70 rotor at 20°C for 3 days. The resulting fluorescent DNA band was removed from the gradient, and treated with isopropanol to extract the ethidium bromide. Finally, the DNA was dialyzed against TE buffer and precipitated by addition of ethanol.

A Nicotiana genomic library was prepared from this DNA as follows, using the phage lambda vector EMBL4 described by Frischauf et al., J. Mol. Bio. 170: 827 (1983). EMBL4 phage was prepared from agarose plate stocks prepared by the method of Davis et al; Advanced Bacterial Genetics, (Cold Spring Harbor Laboratory, New York, 1980). Phage DNA was prepared as described by Silhavy et al., Experiments with Gene Fusions, (Cold Spring Harbor Laboratory, New York, 1984), by concentrating phage with polyethylene glycol, removing the polyethylene glycol by chloroform extraction, and purifying phage using glycerol step gradients. The resulting purified phage was then treated with deoxyribonuclease and ribonuclease prior to phenol extraction. Phage DNA was spooled from ethanol. To prepare arms of the EMBL4 phage, phage DNA was sequentially digested with Sal I and Bam HI endonucleases. The arms were annealed and then separated from the central fragment on a 10-40% sucrose gradient, as described by Maniatis et al., Molecular Cloning: A Laboratory Manual. (Cold Spring Harbor Laboratory, New York, 1982). The arms were completely denatured and reannealed prior to ligation to tobacco DNA. Tobacco DNA, prepared as previously described, was partially digested with Sau3A endonuclease and sedimented through a 10-40% sucrose gradient. Fractions from the sucrose gradient were then analyzed by electrophoresis on 0.5% agarose gels. Fractions containing fragments in the 20-40 kb size range were dialyzed, precipitated, and ligated to the lambda phage DNA arms. The DNA was ligated at a concentration of 135µg per mL vector and 45 µg per mL insert DNA. The resulting ligated concatamers were then packaged using lambda DNA packaging extracts. The resulting yield of phage was approximately 4.5 x 105 phage per μg insert DNA. A library of approximately 400,000 phage was constructed, representing an estimated 99% complete library for tobacco, which has an approximate genomic content of 1.65 picograms, or 1.52 x 10<sup>9</sup> base pairs (Zimmerman, et al., <u>Chromosoma 59:227 1977</u>). The resulting phage library of <u>Nicotiana DNA</u> was grown and plated on <u>E. coll stain LE392 (ATCC 33572)</u>, as

The resulting phage library of Nicotiana DNA was grown and plated on E. coll stain LE392 (ATCC 33572), as disclosed by Silhavy et al., Experiments with Gene Fusions, (Cold Spring Harbor Laboratory, New York, 1984). Phage were plated to provide 2000-5000 plaques on 90 mm petri dishes or 50,000 plaques on 150 mm petri dishes. Plaque lifts were done by the method of Benton et al., Science 196:180 (1977). Following transfer of phage DNA to nitrocellulose filters, the filters were prehybridized by incubation for about 4 hours at 56°C in 6 x SSPE containing 0.5% SDS, 100 μg per mL denatured calf thymus DNA, and 10 x Denhardt's solution. Hybridization was then accomplished as described by Manlatis, et al., Molecular Cloning: A Laboratory Manual, (Cold Spring Harbor Laboratory, New York, 1982) p. 326. In this step, a fresh aliquot of hybridization solution was added, together with about 108 cpm of the radioactive yeast ALS gene probe. Hybridization was allowed to occur for about 24-48 hours at 56°C. At this point, the filters were first rinsed for about 4 hours in 6 x SSPE at 56°C, then rinsed three additional times for 20 minutes each in 2 x SSPE at about 23°C. The filters were then dried and exposed at -70°C for 2 days, using Kodak XAR or XRP x-ray film and a Du Pont Cronex® Lightning Plus® Intensifying screen. Exposed shots on the film indicated the position of plaques potentially containing Nicotiana ALS genes.

The autoradiograms prepared as described above were then oriented over the original bacteriophage-containing petri dishes. Using the wide end of a sterile Pasteur pipette, plaques corresponding to the darkest spots on the autoradiograms were excised. The plaques selected were then eluted into SM buffer and plated onto fresh 90 mm petri dishes. Each dish received about 100-200 phage. The complete phage location process was then reiterated, using freshly prepared probe. In this manner, the phage location and isolation steps were repeated until the majority of plaques indicated the presence of phage containing DNA capable of hybridization to the yeast ALS gene probe.

Mini-preparations of DNA from the plaque-purified phage designation NtA13 were isolated as described and

worked up as described by Maniatis et al. Molecular Cloning: A Laboratory Manual (Cold Spring Harbor Laboratory, New York, 1982), p. 371. EcoRI restriction endonuclease digests of the DNA mini-preparations were electrophoresed through 0.7% agarose gels and blotted onto nitrocellulose filters.

5

15

20

35

40

80

85

Fragments containing the ALS gene were then Identified by hybridization with the yeast ALS gene probe. Fragments capable of hybridization to the probe were then Isolated and subcloned into vectors pBR322, M13mp9, or M13mp18. These fragments were then sequenced using oligonucleotide primers in a dideoxy chain termination procedure conducted substantially as described by Sanger et al., Proc. Natl. Acad. Sci. USA 74:5463 (1977). A kit available from New England Biolabs (Beverly, Massachusetts, U.S.A.) was employed. Use of synthetic oligonucleotide primers allowed extension of a DNA sequence along a cloned fragment in overlapping segments. Computer analysis of the DNA sequence Identified a 667 codon open reading frame. The deduced amino acid sequence of this open reading frame was substantially homologous to the sequences previously determined from the Sacharomyces cerevisiae ILV2 gene and the E. coli ilvG gene, indicating that the DNA fragment recovered from the Nicotiana genomic library contained a tobacco ALS gene. To determine whether this ALS gene encoded the wild type herbicide sensitive enzyme or the mutant herbicide resistant enzyme from the S4 line, the gene was introduced into wild type herbicide sensitive tobacco by Agrobacterium turnefaciens medicated transformation.

A genetic marker for selection of transformed plant cells was required. Resistance to the antibiotic G-418 resulting from expression in plants of a bacterial gene, NPT II, encoding neomycin phosphotransferase was used. To allow expression of NPT II, plant regulatory sequences were fused to the NPT II coding region in the vector pKNK. Vector pKNK was derived from the commonly used plasmid pBR322 by removing the Hind III BamH I sites and inserting at the Cla I site and approximately 2.3 kb Cla I fragment which consisted of the following:

a) a 320 bp Cla I-Bgl II sequence containing the promoter region of the neomycin phosphotransferase (NPT II) gene of transposon Tn 5 derived by the conversion of a Hind III site to the Cla I site [Beck, E., Ludwig, G., Auerswald, E. A., Reiss, B. & Schaller, H. (1982) Gene 19:327-336].

b) a 296 bp Sau 3A-Pst I sequence containing the nopaline synthase promoter (NOSO) derived from the nopaline synthase gene (NOS) (nucleotides -263 to +33, with respect to the transcription start site [Depicker, A., Stachel, S., Dhaese, P., Zambryski, P & Goodman, H. J. (1982) J. Mol. Appl. Genet. 1:561-574] by the creation of a Pst I site at the initiation codon.

c) the 998 bp Hind III- BamH I sequence containing the coding sequence for the NPT II gene derived from Tn 5 by the creation of Hind III and BamH I sites at nucleotides 1540 and 2518 [Beck, E., Ludwig, G., Auerswald, E. A., Reiss, B. & Schaller, H. (1982) Gene 19:327-336], respectively.

d) the 702 bp BamH I-Cla I sequence containing the 3' region of the NOS gene (nucleotides 848 to 1550) [Depicker, A., Stachel, S., Dhaese, P., Zambryski, P. & Goodman, H. J. (1982) J. Mol. Appl. Genet. 1:561-574].

The nucleotide sequence at the fusion of the NOSP and the NPT II coding sequence is:

# NOS Sequence NPT II Sequence ....AATAATCTGCAGCAAGCTTGCGGGGATCGTTCGC ATG .... Pst 1 HindIII

Vector pKNK was cleaved by restriction enzyme Sal I (BRL) and the resultant linear vector, following phenoi extraction, was joined in the presence of T4 DNA ligase (New England Biolabs) to purified 2.1 kb Sal I fragment carrying the bacterial neomyclin phosphotransferase I (NPT I) gene as a selectable marker for bacterial resistance to kanamycin. The ligation mixture was used to transform competent E. coli HB101 cells, and the transformants were selected on plates containing 100 mg/L kanamycin. The resultant recombinant plasmid, designated pKNKS, was purified.

Plasmid pKNKS was cleaved with restriction enzyme <u>Eco</u> RI (BRL), and its ends made blunt by Klenow fragment (BRL). The resultant linear DNA was joined in the presence of T4 DNA ligase (New England Biolabs) to phosphorylated <u>Bgl</u> II linkers. Following extensive digestion with <u>Bgl</u> II restriction enzyme (BRL) to remove the excess linkers, the DNA was passed through a Sephadex G-150 (fine) (Pharmacia) gel filtration column to separate it from the linkers. The DNA fragment was isolated and the volume adjusted to yield a DNA concentration of approximately in the presence of T4 DNA ligase (New England Biolabs) in total volume of 5 µl and used to transform competent <u>E. coli</u> HB101 cells. Ampicillin-resistant cells were shown to contain the plasmid, pKNKSG, which is identical to pKNKS except for the replacement of the <u>Eco</u>RI restriction site with that of Bgl II.

A Sma I fragment of phage NtA13 was shown to contain the region that hybridized to the yeast ALS gene. Phage NtA13 was partially digested by restriction enzyme Sma I (New England Biolabs), and following phenol extraction, joined to phosphorylated BamH I linkers (New England Biolabs) in the presence of T4 DNA ligase (New England Biolabs). Following digestion with BamH I and removal of excess linkers by electrophoresis on an agarose gel, a 16 kb BamH I restriction fragment containing the tobacco ALS gene from Phage NtA13 was isolated from the gel by electroelution and used for further cloning.

Vector pKNKSG was linearized with restriction enzyme Bgl II (BRL) and, following dephosphorylation by calf intestine phosphatase (Boehringer Mannheim), it was joined in the presence of T4 DNA ligase (New England Biolabs) to the 16 kb BamH I restriction fragment derived from phage NtA13. The ligation mixture was used to transform competent E. coli HB101 ceils and an ampicillin-resistant transformant was shown to contain the recombinant plasmid designated pIV13. The orientation of the insert fragment in pIV13 was such that the open reading frames of the NOS:NPT II gene in the vector and the ALS gene in the Insert were in opposite directions. HB101 (pIV13), following purification by three single colony streaking, was used for triparental mating.

Three ml overnight cultures of E. coli HB101 (plV13) and E. coli HB101 (pRK2013) (ATCC number 37159) in LB liquid medium containing 25 mg/L kanamycin were grown at 37°C, and of Agrobacterium tumefaciens GV3850 in LB medium at 28-29°C. The cells were harvested at room temperature in a clinical centrifuge, washed once in LB medium without drug, harvested, and resuspended in 3 ml of LB. 0.25 ml aliquots of all three strains were mixed in a tube and the mixture was transferred onto a Millipore filter (2.5 cm HAWP, 0.45 µm) placed on top of three Whatman No. 1 filters in a petrl dish. After all of the liquid medium was absorbed by the Whatman filter (about 30 min), the Millipore filter with bacteria on its top surface was laid (bacteria side up) onto a LB plate without drug. After incubation overnight at 28-29°C, the Millipore filter was transferred to 5 ml of 10 mM MgSO<sub>4</sub> and vortexed to resuspend the bacteria in the solution. 0.1 ml aliquots were plated on selective plates [M9 minimal plates containing 20% sucrose, 1 mM MgSO<sub>4</sub>, 1 mM CaCl<sub>2</sub>, and 1 mg/ml kanamycin (Sigma)]. Several large colonles showed up after about four days of incubation at 28-29°C. Several transconjugants were purified by three successive single-colony streakings on the same selective plates. Only Agrobacteria containing the plasmid plV13 recombined with the endogenous pGV3850 plasmid through their common pBR322 sequences were expected to grow. This was confirmed by Southern analysis before using the engineered Agrobacterium, GVKNT13, for plant transformations.

For plant cell transformations, standard aseptic techniques for the manipulations of sterile media and axenic plant/bacterial cultures were followed, including the use of a laminar flow hood for all transfers. Recipes for media are given in Example VI. Potted tobacco plants for leaf disk infections were grown in a growth chamber maintained for a 12 hr, 24°C day, 12 hr, 20°C night cycle, with approximately 80% relative humidity, under mixed cool white fluorescent and incandescent lights. Tobacco leaf disk infections were carried out essentially

by the method of Horsch et al. (1985) Science 227, 1229.

Young leaves, not fully expanded and approximately 4-6 inches in length, were harvested with a scalpel from approximately 4-6 week old tobacco plants (Nicotiana tabacum var. Xanthi). The leaves were surface sterilized for 30 minutes by submerging them in approximately 500 ml of a 10% Chlorox, 0.1% SDS solution and then rinsed 3 times with sterile deionized water. Leaf disks, 6 mm in diameter, were prepared from whole leaves using a sterile paper punch.

Leaf disks were inoculated by submerging them for several minutes in 20 ml of a 1:10 dilution of an overnight Agrobacterium culture carrying the plasmid GCKNT13. Agrobacterium cultures were started by inoculating 10 ml of YEB broth with a single bacterial colony removed form a R-agar plate. The culture was grown approximately for approximately 17-20 hours in the 18 mm glass culture tubes in a New Brunswick platform shaker maintained at 28°C.

After inoculation, the leaf disks were placed in petri dishes containing CN agar medium. The dishes were sealed with parafim and incubated under mixed fluorescent and "Gro and Sho" plant light (General Electric) for

2-3 days in a culture room maintained at approximately 25°C.

To rid the leaf disks of <u>Agrobacterium</u> and to select for the growth of transformed tobacco cells, the leaf disks were transferred to fresh CN medium containing 500 mg/L cefotaxime and 100 mg/L kanamycin. Cefotaxime was kept as a frozen 100 mg/ml stock solution and added aseptically (filter sterilized through a 0.45 µm filter) to the media after autoclaving. A fresh kanamycin stock (50 mg/ml) was made for each use and was filter sterilized into the autoclaved media.

Leaf disks were incubated under the growth conditions described above for 3 weeks and then transferred to

fresh media of the same composition.

Approximately 1-2 weeks later, shoots developing on medium containing kanamycin were excised with a sterile scalpel and planted in A medium containing either 10 ppb chlorsulfuron or 100 mg/L kanamycin. Root formation on selective and non-selective media was recorded within 3 weeks.

Within 2 weeks of planting, small leaves were removed from excised shoots to determine levels of resistance to chlorsulfuron and kanamycin in a callus induction assay on selective media. To induce callus formation, small leaves were excised and cut into several sections with a scalpel and planted in B medium containing either 10 ppb chlorsulfuron or 50 mg/L kanamycin. Callus growth on selective and non-selective media was recorded within 3 weeks.

The results shown in Table 1 indicated that transformation of tobacco had been achieved with the GVKNT13 strain based on production of kanamycin resistant callus. The kanamycin resistant callus remained sensitive to the sulfonylurea herbicide chlorsulfuron, indicating that the ALS gene isolated from the tobacco S4 mutant in phage NtA 13 encoded the wild type herbicide sensitive enzyme. This plant ALS gene has been used as a DNA hybridization probe to isolate other plant ALS genes, including genes which encode herbicide resistant ALS.

55

10

30

35

### Table 1

## Results from Callus Tests of GVKNT13\_Infected\_Tobacco

5

40

60

Number of transformed shoot explants producing callus on selective and non-selective media.

		GVKNT13 <sup>1</sup>	<u>gvkk</u> 2	GV3850 <sup>3</sup>	10
Exp	. #1				
Kan	selection amycin, 50 mg/L orsulfuron, 10 ppb	59/62 53/62 0/62	12/13 8/13 0/13	10/10 0/10 0/10	<b>15</b>
Exp	<u>. #2</u>				20
Kan	selection amycin, 50 mg/L orsulfuron, 10 ppb	96/102 81/102 0/102	21/23 16/23 0/23	22/25 0/25 0/25	25
1	Agrobacterium strain tobacco ALS gene and resistance)				<i>30</i>
2	Agrobacterium strain only NOS/NPTII gene	containing 1	Ti plasmid	carrying	
3	Agrobacterium strain of either tobacco AL			(devoid	35

A genomic library of DNA from the Hra mutant of tobacco was made in bacteriophage lambda and screened for clones that hybridized to the wild type tobacco ALS gene from the S4 mutant. Several phage clones were isolated. Physical mapping of the tobacco DNA inserts using restriction endonucleases revealed the presence of two distinct classes of DNA fragments representative of the two tobacco ALS genes SURA and SURB. Comparison of the physical maps of the SURA and SURB genes of N. tabacum to maps from the progenitor species showed that the SURA gene came from N. sylvestris and the SURB gene came from N. tomentosiformis. The wild type ALS gene isolated previously from the S4 mutant was designated SURA. The genetic linkage of the high level herbicide resistance mutation in Hra to the S4 mutation indicated that the Hra mutation was in the same ALS gene as the S4 mutation, namely SURB. Therefore, it was expected that the SURB gene isolated from the Hra mutant would be a mutant gene, designated SURB-Hra, encoding a herbicide resistant ALS. One phage clone containing the SURB-Hra gene was chosen for further analysis. This phage clone, designated 3, has been deposited at the ATCC, Rockville, MD under accession number ATCC 40237. The phage clone was digested with Spe I restriction endonuclease to give an 8.3 Kb DNA fragment which was inserted into the Xba I site of plasmid pMuc19, and the resulting recombinant plasmid, pAGS148, has been deposited at the ATCC, Rockville, MD under accession number ATCC 67124. Plasmids pAGS148 and pAGS135 were ligated to each other as described below, and the resulting recombinant plasmid pAGS152 (Figure 2) was introduced into Agrobacterium tumefaciens LBA 4404. The resultant Agrobacterium tumefaciens LBA 4404 (pAGS152) has been deposited at the ATCC, under accession number ATCC 67126.

A genomic library of DNA from the tobacco C3 mutant was made in bacterlophage lambda and screened for clones which hybridized to the previously isolated ALS genes from tobacco. Several phage clones were isolated and the tobacco DNA inserts where physically mapped with restriction endonucleases. Two different DNA fragment types, corresponding to the SURA-C3 gene and the SURB gene, were identified. Two phage clones designated 35 and 38, carrying the SURA-C3 gene were chosen for further analysis.

Phage clone 35 were digested with Spe I and Sal I restriction endonucleases to give the 6.3 kb DNA fragment shown in Figure 3. This DNA fragment has been inserted into the plasmid vector pUC119 digested

with restriction endonucleases Xba I and Sal I, and the resulting recombinant plasmid, pALS35, has been deposited at the ATCC, Rockville, Maryland, under accession number 67424.

In addition to the four tobacco ALS genes. SURA and SURB encoding wild type herbicide sensitive ALS, and SURA-C3 and SURB-Hra encoding mutant herbicide resistant ALS, ALS genes have been isolated from Arabidopsis thaliana, Beta vulgaris (sugarbeet) and part of an ALS gene from Zea mays (corn). the latter ALS genes, from herbicide sensitive plants, were obtained from genomic DNA libraries made in bacteriophage lambda by screening for DNA hybridizing to a previously isolated ALS gene from yeast or tobacco. The wild type ALS gene from sugarbeet was isolated in a phage designated  $\Phi$ 21 and physically mapped with restriction endonucleases. The DNA fragment isolated in this phage and two DNA fragments which were subcloned into the plasmid vector pUC119 are shown in Figure 7. Plasmid pSBALS216 has been deposited at the ATCC, Rockville, Maryland under accession number 67425.

10

30

35

40

45

50

55

60

65

Figure 1 shows restriction endonuclease maps of DNA fragments containing ALS genes isolated from the Hra mutant of tobacco. Based on these maps, two classes of DNA fragments can be distinguished. An approximately 18 kilobase nucleic acid insert in phage clone 3 carries the SURB-Hra gene. The insert contains a preferred DNA fragment of the present invention which encodes a herbicide-resistant ALS from tobacco mutant Hra. This nucleic acid fragment consists of double-stranded DNA of 8.3 ± .5 kilobases and has a molecular weight of 5.5 ± .3 mega daltons, and has 5' overhang sequences of CTAG at both ends. The 8.3 kilobase nucleic acid fragment between the two Spe I sites hybridized to the ALS gene probe used to screen the genomic library. Restriction endonuclease Spe I can be used to excise the fragment from the phage using well-known techniques.

Figure 2 shows a physical map of plasmid pAGS152. Plasmids pAGS135 and pAGS148 are not drawn to scale. Restriction endonuclease sites EcoR I (RI), BamH I (B), Xba I (X), Pst I (P), Sal I (S), Spe I (Sp), Nco I (N), Hind III (H), BstE II (Bs), Sma I (Sm), Kpn I (K), Sst I (St), Sph I (Sh) and Bgl II (G) are shown. pAGS152 results from the ligation of the BamH I- cleaved plasmids pAGS135 (approximately 27 kilobases) and pAGS148 (approximately 12.1 kilobases). Plasmid pAGS135, drawn as a circle, is a wide host range plasmid containing a plant kanamycin resistance gene (NOS:NPT II) and a BamH I cloning site, flanked by the left border (LB) and the right border (RB) of T-DNA. Plasmid pAGS148, shown as a linear BamH I fragment, consists of the Spe I (Sp) fragment (approximately 8.3 kilobases) of the aspect of the invention (shown flanked by X/Sp and Sp/X), containing the coding sequence for the herbicide-resistant form of ALS, from the Hra mutant, inserted in the Xba I site (X) of plasmid pMuc19 (open box). Although Spe I and Xba I restriction enzymes recognize different sequences, their action results in DNA fragments with the same 5' overhanging sequence, viz 5'-CTAG-3'. Thus, Spe I and Xba I digest dfragments can be ligated to each other, but the ligation results in a loss of both sides. The hatched box on the insert fragment corresponds to the coding region of the ALS gene and the arrow denotes the 5'-3' direction of the coding sequence. The nucleic acid fragment is flanked by Hind III, Sph I, Pst I and Sal I sites at one end and by BamH I, Sma I, Kpn I, Sst I and EcoR I sites at the other end. These enzymes can be used to excise the fragment from the plasmid by complete or partial digestion using well-known techniques. After digestion, the ends of the fragment will be characteristic of the endonuclease used to excise the fragment:

- - - - - -

		0 257	993		
•	5' Overha	inging Sequence	3' Ove	rhanging Sequence	
	Spe I	5'-CTAGT-3' 3'-A-5'	Sph I	5'-C-3' 3'-GTACG-5'	5
	: <u>Hin</u> d III	5'-AGCTT-3' 3'-A-5'	Pst I	5'-G-3' 3'-ACGTC-5'	10
	Sal I	5'-TCGAC-3' 3'-G-5'	<u>Kpn</u> I	5'-C-3' 3'-CATGG-5'	
	<u>Bam</u> H I	5'-GATCC-3' 3'-G-5'	<u>Sst</u> I	5'-C-3' 3'-TCGAG-5'	
	EcoR I	5'-AATTC-3' 3'-G-5'			20
	<u>Sma</u> I	Blunt end 5'-GGG-3' 3'-CCC-5'			25
fragmen ALS from	it can be characteriz m mutant plant Hra	zed by the restriction map sl of Nicotiana tabacum cv. "	hown in Figu Kanthi' which	st using agarose gel electrophoresis. The re 2, and contains the coding sequence for a resistant to inhibition by chlorsulfuron ucleotide sequences required to express	30
the gen Figure which co with res	e in plants. e 3 shows a restrict arries the <u>SURA-C3</u> triction endonuclea	ion endonuclease map of th gene. This DNA fragment v ses Spel and Sall and was in	e approxima vas obtained serted into t	tely 6.8 kb preferred nucleic acid fragment from lambda phage clone 35 by digestion he plasmid vector pUC119 which had been	35
digested with restriction endonucleases Xba I and Sal I, as described in the legend to Figure 2.  Figure 4 shows a partial nucleotide sequence of a preferred DNA fragment encoding a herbicide-resistant form of ALS from SURB-Hra gene of tobacco. Nucleotides are indicated by their bases by the following standard abbreviations:  A = adenine;					
C = cyt T = thy G = gu	mine; anine.				45
The begining of the nucleotide sequence corresponds to Pst I Site (P) 885 nucleotide bases preceding the coding sequence, shown on Figure 2; the sequence ends at base number 2946, which is 67 bases past the end of the coding sequence shown in Figure 2. The nucleotide sequence from nucleotide one to nucleotide 884 is believed to contain 5' regulatory sequence(s) required for expression of the encoded ALS. Figure 4 also shows the deduced amino acid sequence of the ALS protein.  Amino acid residues are indicated by the following abbreviations:					
A = alanine; C = cysteine; D = aspartic acid; E = characteristics					

E = glutamic acid; F = phenylalanine;

G = glycine; H = histidine;

! = isoleucine; K = lysine;

L - leucine;

M = methionine;

N = asparagine;
P = proline;

11

65

Q = glutamine;

R = arginine;

S = serine:

T = threonine;

V - valine;

20

25

35

40

50

55

W = tryptophan; and

Y = tyrosine.

The term "amino acids" as used herein is meant to denote the above-recited natural amino acids and functional equivalents thereof.

Figure 5 shows a partial nucleotide sequence and its cognate deduced amino acid sequence, of a preferred DNA fragment encoding a herbicide-resistant form of ALS from the C3 gene of tobacco. The beginning of the nucleotide sequence corresponds to the BamH I site shown in Figure 3. The coding sequence begins at nucleotide 176 and ends at nucleotide 2175. The nucleotide sequence from nucleotide one to nucleotide 175 is believed to contain 5' regulatory sequence(s) necessary, but not sufficient, for expression of the encoded ALS. Nucleotides and amino acids are indicated by the standard abbreviations, as shown above.

Figure 6 shows the deduced amino acid sequences of the large subunits of ALS Isozymes I, II and III from E. coll (Lines E, F and G respectively), wild type ALS proteins of yeast (Line D), Arabidopsis thaliana (Line C) and Nicotiana tabacum (tobacco) (Lines A and B), encoded by the SURB and SURA genes, respectively. Amino acid residues are indicated by standard abbreviations as shown above. The first amino acid, methionine, of the deduced amino acid sequences of the yeast (line D, Figure 6) and higher plant (lines A-C) ALS proteins is the putative start of the transit peptides believed to be involved in translocating the enzymes into mitochondria, in the case of the yeast enzyme, or chloroplasts, in the case of the plant enzymes. These transit peptides are believed to be cleaved off during translocation of the proteins into the organelles and are believed not to be required for ALS activity. The extend of these transit peptides is difficult to determine in the absence of data on the In vivo N-termini of the ALS proteins of yeast and higher plants. Based on the homology with the bacterial ALS proteins the chloroplast and mitochondrial transit sequences may be estimated to extend for 90 amino acids.

The dotted lines in the sequences are spacing marks inserted to best align regions of homology. Vertical lines highlight the amino acid residues that are conserved between adjacent sequences of Figure 6. The homology between tobacco and <u>Arabidopsis</u> ALS proteins (lines A to C), which derive from two different plant families, is striking. Even more unexpected, considering the evolutionary distance between microbes and higher plants, is the finding that the amino acid residues which are conserved between the bacterial (lines E to G) and the yeast (line D) ALS proteins are largely conserved between these proteins and the plant ALS proteins.

Figure 7 shows a restriction endonuclease map of the approximately 17.5 kilobase nucleic acid insert in phage clone  $\phi$ 21 carrying the sugarbeet ALS gene. Two smaller DNA fragments which also contain the sugarbeet ALS gene and which were subcloned into the pUC119 plasmid vector are also shown.

Figure 8 shows the deduced amino acid sequences of wild type ALS proteins from the plants Nicotiana tobacum (tobacco) (Lines A and B), Arabidopsis thaliana (Line C) Beta vulgaris cv. sennica (sugarbeet) (Line D) and a partial sequence of the ALS protein from maize (Line E). The dotted lines in the sequences are spacing marks to best align regions of homology. Vertical lines highlight the amino acid sequences that are conserved between adjacent sequences. The homology between all of the plant ALS proteins is very extensive. Based upon this, a mutation in one plant ALS gene causing an amino acid substitution that results in sulfonyturea herbicide resistant ALS would be expected to have an analogous effect if it were present in any other plant ALS gene

The amino acid residues which are conserved in all of the ALS sequences in Figure 6 are believed to be important for the binding of substrates, herbicides, coenzymes, etc. These sequences are believed to be substantially conserved in all ALS proteins. The residues which are partially conserved in the different ALS proteins may participate in less conserved aspects of enzyme function, such as those which govern its herbicide sensitivity and its end-product inhibition. Examples of this would include the resistance of bacterial isozyme I to sulfometuron methyl and chlorsulfuron, and of bacterial isozyme II to end-product inhibition by valine. Finally, those residues which are not conserved between the proteins probably reside in the framework of the ALS protein where sequence divergence is less disruptive to enzyme function.

Although not wishing to be bound by theory, binding of sulfonylurea herbicides to ALS during acetolactate synthesis is believed to be facilitated by the binding of a first pyruvate molecule to the enzyme. However, the binding of a sulfonylurea herbicide molecule is competitive with the binding of a second pyruvate molecule to the enzyme. Sulfonylurea herbicide sensitivity is conserved through evolution in most ALS enzymes. From these facts, it was deduced that the binding of the sulfonylurea herbicide occurs at or proximal to one or more of the conserved amino acids in the ALS proteins. In fact, Applicant has discovered that substitutions for one or more of 10 specific amino acid residues in one or more of the 7 substantially conserved sub-sequences A through G will confer herbicide residence and are claimed. It is expected that substitution at other amino acid residues in the substantially conserved sub-sequences will also confer herbicide resistance.

Sulfonylurea herbicide resistance in bacteria, yeast and higher plants, which resistance cosegregates with herbicide-resistant forms of ALS, results from mutations in the structural genes for ALS. Comparing the nucleotide sequences of ALS genes of organisms encoding herbicide sensitive and herbicide-resistant forms

of ALS allows one to determine which amino acid residues are important for herbicide inhibition of the enzyme. One mutation in the <u>E. coli ivG</u> gene, which results in an enzyme with increased resistance to suffometuron methyl inhibition, and with reduced catalytic activity, was determined to result in an alanine-to-valine substitution at position 122 (Figure 6). Another sulfometuron methyl resistance mutation in this gene was determined to result in a alanine-to-serine substitution at the same position. This alanine residue is conserved in all ALS enzymes except bacterial isozyme I (Figure 6), which is naturally resistant.

Many genes encoding herbicide-resistant ALS enzymes have been isolated from spontaneous sulfonylurea-resistant yeast mutants. Sequencing of these genes has shown the molecular basis of resistance to be base changes which result in amino acid substitutions at ten different positions in the protein (Table 2), residues 121, 122, 197, 205, 256, 359, 384, 588, 591 and 595 (numbering relative to the positions in Figure 6).

0 257 993

Table 2

Spontaneous Mutations of the Yeast ALS Gene Resulting in Sulfonylurea Herbicide Resistance

5					
	Amino Acid Positions	Wild Type Codon	Wild Type Amino Acid	Mutant Codon	Amino Acid Substitution
10	121	GGT	Gly	<u>A</u> GT	Ser
	122	GCT	Ala	<u>C</u> CT G <u>A</u> T	Pro Asp
15				G <u>T</u> T <u>A</u> CT	Val Thr
	197	CCA	Pro	<u>T</u> CA C <u>G</u> A	Ser Arg
20	205	GCT ·	Ala	G <u>A</u> T <u>A</u> CT	Asp Thr
25	256	AAG	Lys	<u>G</u> AG A <u>C</u> G AA <u>C</u>	Glu Thr Asn
	359	ATG	Met	<u>G</u> TG	Val
30	384	GAC	Asp	GA <u>A</u> G <u>T</u> C <u>A</u> AC	Glu Val Asn
35	588	GTT	Val	G <u>C</u> T	Ala
	591	TGG	Trp	<u>C</u> GG	Arg Arg
40				TG <u>T</u> TG <u>C</u> <u>G</u> GG	Cys Cys Gly
45				TIG TCG GCG	Leu Ser Ala
•	595	TTC	Phe	TT <u>A</u>	Leu

*50* 

*55* 

At six of these positions, 122, 197, 205, 256, 384 and 591 (Table 2), more than one substitution that confers herbicide resistance has been obtained. At position 122, at which an alanine residue is present in all known

wild type ALS enzymes except <u>E. coli</u> isozyme I, substitutions of aspartic acid, proline, threonine or valine result in sulfonylurea-resistant ALS. At position 197, at which a proline residue is present in all known wild-type ALS enzymes except <u>E. coli</u> isozymes II and III, substitutions of serine or arginine result in sulfonylurea-resistant ALS. At position 205, at which an alanine residue is present in all known wild type ALS enzymes, substitutions of aspartic acid of threonine result in sulfonylurea-resistant ALS. At position 256, at which a lysine residue is present in all known wild type ALS enzymes, substitutions of glutamic acid, asparagine or threonine result in sulfonylurea-resistant ALS. At position 384 at which an aspartic acid is present in all known wild type ALS enzymes, substitutions of glutamic acid, asparagine or valine result in sulfonylurea-resistant ALS. At position 591, at which a tryptophan is present in all known wild type ALS enzymes except <u>E. coli</u> isozyme I, substitutions of alanine, cysteine, glycine, leucine, arginine or serine result in sulfonylurea-resistant ALS.

Mutants resistant to sulfonylurea herbicides resulting from single amino acid substitutions at the other four positions, 121, 359, 588, 595, have been obtained. At position 121, at which glycine is present in all known ALS enzymes, substitution of serine results in a sulfonylurea-resistant ALS. At position 359, at which methionine is present in all known ALS enzymes, substitition of valine results in a sulfonylurea-resistant ALS. At position 588, at which valine is present in all known ALS enzymes, substitution of alanine results in sulfonylurea-resistant ALS. At position 595, at which phenylalanine is present in all known ALS enzymes except E. coli isozyme ill, substitution of leucine results in sulfonylurea-resistant ALS.

Oligonucleotide-directed site specific mutations, which result in amino acid substitutions at positions 122, 205, 256, 359, 384 and 591, have been made in the yeast gene encoding ALS (Table 3).

Table 3

Site-Directed Mutations of the Yeast ALS Gene Resulting in Sulfonylurea Herbicide Resistance

Amino Acid Positions	Wild Type Codon	Wild Type Amino Acid	Mutant Codon	Amino Acid Substitution	0.
122	GCT	Ala	<u>TC</u> T G <u>T</u> T <u>A</u> CT <u>C</u> CT	Ser Val Thr Pro	30
			AAT ATT CAT CGT	Asn Ile His Arg	35
			CIT TAT TGT TTT	Leu Tyr Cys Phe	40
			GAA ATG AAA CAA TGG	Glu Met Lys Gln Trp	45
205	GCT	Ala	CGT TGT GAA TGG	Arg Cys Glu Trp	50 55
<b>256</b> .	AAG	Lys	GAC GGG CTG CCG TGG	Asp Gly Leu Pro Trp	60

10

15

20

0 257 993

## Table 3 (continued)

5	Amino Acid Positions	Wild Type Codon	Wild Type Amino Acid	Mutant Codon	Amino Acid Substitution
3	359	ATG	Met	CCA CAG	Pro Glu
10 15	384	GAC	Asp	CCA TGG TCC GGT TGC AAA	Pro Trp Ser Gly Cys Lys
20	591	TGG	Trp	GAC GAG TTC CAC TAC ATA	Asp Glu Phe His Tyr Ile
25	ē	<i></i>		<u>GT</u> G <u>AA</u> G <u>A</u> GG <u>AT</u> G	Val Lys Arg Met Asn
30			1	AAC CAG ACG	Gln Thr

35

At position 122, mutations resulting in eighteen amino acid substitutions for alanine, which is present in wild type ALS, have been made. The nineteenth substitution (aspartic acid) was isolated previously as a spontaneous mutation and was therefore not remade. Each substitution, except for glycine, results in sulfonylurea-resistant ALS. At position 205, mutations resulting in substitutions for alanine, the wild type residue, of cysteine, glutamic acid, arginine or tryptophan, result in sulfonylurea-resistant ALS. At position 256, mutations resulting in substitutions for tysine, the wild type residue, of aspartic acid, glycine, leucine, proline or tryptophan, result in sulfonylurea-resistant ALS. At position 359, mutations resulting in substitutions for methionine, the wild type residue, of glutamic acid or proline result in sulfonylurea-resistant ALS. At position 384, mutations resulting in amino acid substitutions for aspartic acid, the wild type residue, of cysteine, glycine, proline, lysine, serine or tryptophan result in sulfonylurea-resistant ALS. At position 591, mutations resulting in amino acid substitutions for tryptophan, the wild type residue, of aspartic acid, glutamic acid, phenylalanine, histidine, isoleucine, lysine, arginine, valine, methionine, asparagine, glutamine, threonine or tyrosine result in sulfonylurea-resistant ALS.

All mutations described in Tables 2 and 3 resulted in enzymes which were active and less inhibited by sulfonylurea herbicides than the wild type. Taken in total these results indicate that most substitutions at these 10 positions result in enzymatically active herbicide resistant ALS.

The deduced amino acid sequences of the wild type ALS proteins from tobacco, Arabidopsis, sugarbeet and corn are shown in Figure 7. The amino acid residues at positions 121, 122, 197, 205, 256, 359, 384, 588, 591 and 595 (numbering of positions from Figure 6) in all the plant enzymes are the same as those present in the wild type herbicide sensitive yeast protein (Figure 6). The deduced amino acid sequence of the tobacco ALS gene SURB-Hra, which encodes a herbicide-resistant ALS, is shown in Figure 4. The mutant gene of Figure 4 was derived from a tissue culture line which had undergone two successive spontaneous mutations. The two mutations have been shown to be genetically linked, and introduction of this fragment into sensitive tobacco cells confers upon the cells the same level of herbicide resistance as is found for the original highly resistant mutant tobacco plant from which the fragment was derived. Based on these facts, it was expected that there would be two amino acid substitutions in the enzyme encoded by the fragment. A comparison of the deduced amino acid sequence of the mutant ALS with the deduced amino acid sequence of the wild type ALS reveals that the mutant ALS has a proline-to-alanine substitution at position 197 (Figure 6) and a tryptophan-to-leucine substitution at position 591 (Figure 6). Based on the foregoing, it was determine that substitutions at proline 197 and tryptophan 591 residues confer herbicide resistance. The deduced amino acid sequence of a second

mutant tobacco ALS gene, <u>SURA-C3</u>, which encodes a sulfonylurea herbicide resistant ALS, is shown in Figure 5. A comparison of the deduced amino acid sequences of the mutant and wild type ALS enzymes (Figure 5 and Figure 6, line B) reveals that the mutant ALS has a single substitution, proline-to-glutamine, at position 197. The C3 cell line from which the <u>SURA-C3</u> gene was obtained showed selective herbicide resistance. That is, the C3 mutation conferred resistance to the sulfonylurea herbicides chlorsulfuron and sulfomenturon methyl, but not to an imidazolinone herbicide.

The identification of amino acid substitutions in herbloide-resistant ALS enzymes from plants at positions 197, from the C3 and Hra mutants, and 591 from the Hra mutant, indicates that substitutions at positions

operable in yeast ALS are also operable in plant ALS.

While the amino acid residues present at positions, 121, 122, 197, 205, 256, 359, 384, 588, 591 and 595 are conserved in all wild type herbicide sensitive ALS enzymes so far characterized from eucaryotes, some substitutions at these positions are found in wild type bacterial ALS enzymes. E. coli isozyme I has a serine rather than alanine at position 122 and a glutamine rather than tryptophan at position 591, E. coli isozyme II has a serine rather than proline at position 197 and E. coli isozyme III has an alanine rather than proline at position 197 and an isoleucine rather than phenylalanine at position 595. Each of these E. coli ALS isozymes is more resistant (from 50-fold to greater than 10,000-fold) to inhibition by (particular) sulfonylurea herbicides than plant or yeast ALS. Furthermore, a site-directed mutation causing a serine-to-proline substitution at position 197 in E. coli ALS II rendered the mutant enzyme 100 fold more sensitive to inhibition, i.e., as sensitive as wild type higher plant enzymes. Thus, proline at position 197 is involved in herbicide binding in E. coli ALS II as well as in yeast and higher plant ALS.

15

30

In addition, site-directed mutations which result in tryptophan-to-leucine and glutamine-to-tryptophan substitutions at position 591 in ALS II and ALS I, respectively, of <u>E. coli</u> have been made. The mutation in ALS II makes the enzyme more herbicide resistant than the wild type ALS II, while the mutation in ALS I makes it more

sensitive than wild type ALS I.

The site-directed mutations at positions 197 and 591 in ALS I and ALS II of <u>E. coli</u> affect inhibition by herbicide of the mutant enzymes in a manner predicted from the herbicide-resistant mutant yeast and plant ALS proteins. These experimental findings support the universality of the amino acid residues involved in herbicide binding to ALS enzymes from diverse sources.

Characterization of Nucleic Acid Fragments Encoding Herbicide-Resistant ALS

According to the present invention, the amino acid residues of ALS that correspond to  $\epsilon_1$  and  $\epsilon_2$  in amino acid sub-sequence A,  $\alpha_1$  in amino acid sub-sequence B,  $\delta_2$  in amino acid sub-sequence C,  $\lambda_1$  in amino acid sub-sequence D,  $\gamma_1$  in amino acid sub-sequence E,  $\beta_3$ ,  $\beta_7$  and  $\beta_8$  in amino acid sub-sequence F and  $\sigma_1$  in amino acid sub-sequence G of Figure 6 (referred to hereinafter as positions 122, 121, 197, 205, 256, 384, 591, 595, 588 and 359 respectively) are important in herbicide sensitivity or resistance of ALS enzymes regardless of the biological source of these enzymes, and any nucleotide sequence encoding a plant ALS can be altered to direct synthesis of a herbicide-resistant ALS by virtue of amino acid substitutions at these residues. The nucleic acid fragment of the present invention is characterized in that at least one of the following conditions is met:

a) The nucleic acid fragment encodes an amino acid other than glycine at position 121. Preferably the amino acid is serine, threonine, or cysteine. Most preferably the amino acid is serine.

b) The nucleic acid fragment encodes an amino acid other than alanine at position 122. Most preferably, the amino acid is any other than glycine.

c) The nucleic acid fragment encodes an amino acid other than proline at position 197. Preferably, the amino acid is alanine, glycine, arginine, lysine, histidine, serine, threonine, cysteine, glutamine, or asparagine. Most preferably, the amino acid is alanine, arginine, serine or glutamine.

d) The nucleic acid fragment encodes an amino acid other than alanine at position 205. Preferably, the amino acid is threonine, serine, cysteine, tyrosine, aspartic acid, glutamic acid, tryptophan, histidine, phenylalanine, arginine or lysine. Most preferably, the amino acid is threonine, cysteine, aspartic acid, glutamic acid, tryptophan or arginine.

e) The nucleic acid fragment encodes an amino acid other than lysine at position 256. Preferably, the amino acid is glycine, alanine, leucine, isoleucine, valine, threonine, serine, cystelne, tryosine, glutamic acid, aspartic acid, proline, asparagine, glutamine, tryptophan or phenylalanine. Most preferably, the amino acid is glycine, leucine, threonine, glutamic acid, aspartic acid, proline, asparagine or tryptophan.

f) The nucleic acid fragment encodes an amino acid other than methionine at position 359. Preferably, the amino acid is glutamic acid, aspartic acid, proline, valine, leucine or isoleucine. Most preferably the amino acid is glutamic acid, proline or valine.

g) The nucleic acid fragment encodes an amino acid other than aspartic acid at position 384. Preferably, the amino acid is glycine, alanine, valine, leucine, isoleucine, serine, threonine, cysteine, tyrosine, glutamic acid, proline, asparagine, glutamine, lysine, arginine, tryptophan, histidine or phenylalanine. Most preferably, the amino acid is glycine, valine, serine, cysteine, glutamic acid, proline, asparagine, lysine or tryptophan.

h) The nucleic acid fragment encodes an amino acid other than valine at position 588. Preferably the amino acid is alanine or glycine. Most preferably the amino acid is alanine.

i) The nucleic acid fragment encodes an amino acid other than tryptophan at position 591. Most

preferably, the amino acid is other than proline.

j) The nucleic acid fragment encodes an amino acid other than phenylalanine at position 595. Preferably, the amino acid is leucine, isoleucine and valine. Most preferably the amino acid is leucine.

In one embodiment, position 121 resides within amino acid sub-sequence A as follows:

5 PGε<sub>2</sub>Α

wherein P, G and A are as defined above. To confer herbicide resistance  $\epsilon_2$  is an amino acid other than glycine. Most preferably  $\epsilon_2$  is the amino acid serine. This sub-sequence begins about 24 residues from the begining of a substantially conserved amino acid sequence HEQ.

10 i.e.,

PG<sub>62</sub>A...HEQ.

In one embodiment, position 122 resides within amino acid sub-sequence A as follows:

PGGe:

wherein P and G are as defined above. To confer herbicide resistance ε<sub>1</sub> is a natural amino acid other than alanine. Most preferably ε<sub>1</sub> is any amino acid except glycine. This sub-sequence begins about 24 residues from the begining of a substantially conserved amino acid sequence HEQ,

i.e.,

PGGE1...HEQ.

20 In one embodiment, position 197 resides within amino acid sub-sequence B as follows:

wherein G, Q, and V are as defined above, to confer herbicide resistance  $\alpha_1$  is an amino acid other than proline. Most preferably,  $\alpha_1$  is alanine, arginine, serine or glutamine. This sub-sequence, begins about 20 residues from the end of one substantially conserved amino acid sequence

and about 55 residues from the beginning of a second substantially conserved amino acid sequence SGRPGP.

i.e.,

30

SGPGATN...GQVa1....SGRPGP.

In one embodiment, position 205 resides within an amino acid sub-sequence C as follows:

IGδ<sub>1</sub>Dδ<sub>2</sub>FQE

wherein I, G, D, F, Q, and E are as defined above,  $\delta_1$  represents an amino acid residue which can vary according to the source of the enzyme, but is most commonly T. To confer herbicide resistance  $\delta_2$  is an amino acid other than alanine. Most preferably,  $\delta_2$  is threonine, cysteine, aspartic acid, glutamic acid, arginine, or tryptophan. This sub-sequence begins about 5 residues from the end of a substantially conserved amino acid sequence

GOV

and about 43 residues from the begining of a second substantially conserved amino acid sequence SGRPGP.

*io* i.e.

GQV...IG81D82FQE...SGRPGP.

In one embodiment, position 256 resides within an amino acid sub-sequence D as follows:

Ž1D

wherein P and D are as defined above. To confer herbicide resistance  $\lambda_1$  is an amino acid other than lysine. Preferably,  $\lambda_1$ , is glycine, leucine, threonine, glutamic acid, aspartic acid, asparagine, tryptophan or proline. This sub-sequence D begins about 6 residues from the end of a substantially conserved amino acid sequence SGRPGP

i.e.,

50

SGRPGP...pλ<sub>1</sub>D.

In one embodiment, position 359 resides within an amino acid sub-sequence G as follows

wherein M, L, G and H are defined as above. To confer herbicide resistance,  $\sigma_1$  is an amino acid other than methionine. Most preferably,  $\sigma_1$  is proline, glutamic acid or valine. This sub-sequence ends about 20 residues from the beginning of a substantially conserved amino acid sequence

55 RFDDR

i.e.,

MLGo1HG...RFDDR.

In one embodiment, position 384 resides within an amino acid sub-sequence E as follows

wherein R, F, and D are as defined above. To confer herbicide resistance, γ<sub>1</sub> is an amino acid other than aspartic acid. Most preferably, γ<sub>1</sub> is glycine, valine, cysteine, serine, lysine, glutamic acid, proline, asparagine or tryotophan.

This sub-sequence begins about 20 residues from the end of a substantially conserved amino acid sequence

65 MGLMHG,

l.e., MLGMHGRFD <sub>Y1</sub> R.	
In one embodiment, position 588 resides within an amino acid sub-sequence F as follows	
Gβ1βεβ2Qβ3β4β5β5β7 wherein G and Q are defined above, β1 to βε will vary depending upon the source of the enzyme. β1 is usually	5
methionine, $\beta_3$ is usually tryptophan and $\beta_7$ is usually phenylalanine. To confer herbicide resistance $\beta_8$ is an amino acid other than valine. Most preferably $\beta_8$ is alanine. This sub-sequence begins about 49 residues from the end of a substantially conserved amino acid sequence	
GLPAA	
i.e., GLPAAGβ1β8β2Qβ3β4β5β6β7.	10
In one embodiment, position 591 resides within an amino acid sub-sequence F as follows Gβ <sub>1</sub> Vβ <sub>2</sub> Qβ <sub>3</sub> β <sub>4</sub> β <sub>5</sub> β <sub>6</sub> β <sub>7</sub>	
wherein, G, V and Q are defined above, $\beta_1$ to $\beta_7$ will vary depending upon the source of the enzyme. $\beta_1$ is usually methionine and $\beta_7$ is usually phenyalanine. To confer herbicide resistance $\beta_3$ is any amino acid other than tryptophan. Most preferably $\beta_3$ is any amino acid other than proline. This sub-sequence begins about 49 residues from the end of another substantially conserved amino acid sequence GLPAA.	15
i.e.,	
GLPAAGβ <sub>1</sub> Vβ <sub>2</sub> qβ <sub>3</sub> β <sub>4</sub> β <sub>5</sub> β <sub>5</sub> β <sub>7</sub> . In one embodiment, position 595 resides within an amino acid sub-sequence F as follows Gβ <sub>1</sub> Vβ <sub>2</sub> Qβ <sub>3</sub> β <sub>4</sub> β <sub>5</sub> β <sub>6</sub> β <sub>7</sub>	20
wherein G. V and Q are defined above. $\beta_1$ to $\beta_7$ will vary depending upon the source of the enzyme. $\beta_1$ is usually methionine and $\beta_3$ is usually tryptophan. To confer herbicide resistance, $\beta_7$ is an amino acid other than	
phenylalanine. Most preferably, β <sub>7</sub> is leucine. This sub-sequence begins about 49 amino acids from the end of a substantially conserved amino acid sequence GLPAA	25
i.e.,	
GLPAAGβ <sub>1</sub> Vβ <sub>2</sub> Qβ <sub>3</sub> β <sub>4</sub> β <sub>5</sub> β <sub>5</sub> β <sub>7</sub> .  Herbicide resistance can be achieved by any one of the above described amino acid substitutions and by combinations thereof.	30
The precise amino acid substitutions required for herbicide resistance can be achieved by mutating a	
nucleic acid fragment encoding a herbicide sensitive ALS from any plant of interest generally as follows:  (1) isolate genomic DNA or mRNA from the plant;	
<ul><li>(2) prepare a genomic library from the isolated DNA or a cDNA library from the isolated RNA;</li><li>(3) identify those phages or plasmids which contain a DNA fragment encoding ALS;</li></ul>	35
<ul><li>(4) sequence the fragment encoding the ALS;</li><li>(5) sub-clone the DNA fragment carrying the ALS gene into a cloning vehicle which is capable of</li></ul>	
producing single-stranded DNA;	
(6) synthesize an oligonucleotide of about 15 to 20 nucleotides which is complementary to a particular ALS nucleotide sequence encoding one of the amino acid sub-sequences recited above except for the nucleotide change(s) required to direct a mutation to a codon for an amino acid selected for its ability to	40
confer herbicide resistance; (7) anneal the oligonucleotide to the single-stranded DNA containing the region to be mutated and use it to prime synthesis in vitro of a complementary DNA strand forming a heteroduplex;	45
<ul> <li>(8) transform bacterial cells with the heteroduplex DNA;</li> <li>(9) screen the transformed bacterial cells for those cells which contain the mutated DNA fragment by a) immobilizing the DNA on a nitrocellulose filter, b) hybridizing it to the 5'-32P labelled mutagenic</li> </ul>	
oligonucleotide at ambient temperature, and c) washing it under conditions of increasing temperature so as to selectively dissociate the probe from the wild-type gene but not the mutant gene;  (10) isolate a double-stranded DNA fragment containing the mutant gene; and	50
(11) confirm the presence of the mutation by DNA sequence analysis.  An amino acid substitution required for herbicide resistance can also be achieved by substituting a	
nucleotide sequence of a plant ALS gene which encodes a sequence of amino acids containing the amino acid to be substituted with another nucleotide sequence, which encodes the corresponding stretch of amino acids containing the desired substitution, derived from any natural ALS gene (including microbial) or from a synthetic source.	55
Preparation of Herbicide-Resistant Plants  The public self fragments of the present invention can be used to introduce berbicide registeres into	~~
The nucleic acid fragments of the present invention can be used to introduce herbicide resistance into plants. In order to introduce a nucleic acid fragment which includes a gene encoding herbicide resistant ALS into different plants, a wide variety of techniques are used depending on the species or cultivar desired. In general, explants or protoplasts can be taken or produced from either in vitro or soil grown plants. Explants or protoplasts may be produced from cotyledons, stems, petioles, leaves, roots, immature embryos, hypocotyls,	60
inflorescences, etc. In theory, any tissue which can be manipulated in vitro to give rise to new callus or	65

organized tissue growth can be used for genetic transformation.

5

20

25

30

To achieve transformation, explants or protoplasts may be cocultured with Agrobacterium, which can be induced to transfer nucleic acid fragments located between the T-DNA borders of the TI plasmid to the plant cells. Another method, less commonly used, is direct DNA uptake by plant protoplasts. With this method, the use of Agrobacterium is bypassed and DNA is taken up directly by the protoplasts under the appropriate conditions.

In the examples, a variety of explants from different plants have been cocultured with Agrobacterium to achieve transformation to herbicide resistance. These explants were cultured to permit callus growth. The callus was then tested directly for resistance to sulfonylureas, or plants were regenerated and the plants were tested for sulfonylurea resistance. Testing consisted of an enzyme assay of plant cell extracts for the presence of ALS activity resistant to herbicide and/or growth of plant cells in culture or of whole plants in the presence of normally inhibitory concentrations of herbicide.

The DNA fragments are comprised of a region coding for the synthesis of herbicide-resistant ALS and a region providing for expression of the coding sequence in plants. The 8.3 kb DNA fragment shown in Figure 2 which codes for the herbicide-resistant ALS protein shown in Figure 4 contains about 800 bp in the 5' direction (upstream) of the coding region, sufficient for expression of the protein in plants. This DNA fragment can confer resistance to chlorsulfuron up to 2000 ppb in transformed tobacco calluses. Plants regenerated from the transformed cells also show resistance at the whole plant level. The 6.3 kb DNA fragment shown in Figure 3 which codes for the herbicide resistant ALS protein shown in Figure 5 contains 2.5 kb in the 5' direction (upstream) and 1.8 kb in the 3' direction (downstream) of the coding region sufficient for expression of the protein in plants. This DNA fragment can confer resistance to chlorsulfuron at 2ppb in transformed tobacco

In work which is on-going, DNA fragments containing site-directed mutations in the SURA gene that are expected to code for herbicide resistant ALS have been made. These mutations result in the following amino acid substitutions: Ala 122 to Ser, known to be operable in E. coli ALS Isozyme II and yeast ALS, Ala 122 to Val, known to be operable in E. coli ALS isozyme II and yeast ALS, Ala 122 to Pro, known to be operable in yeast ALS, Pro 197 to Ser, known to be operable in yeast ALS, and E. coli ALS II enzyme, Pro 197 to Ala, known to be operable in ALS encoded by the SURB-Hra gene of tobacco, Lys 256 to Glu, known to be operable in yeast ALS, Asp 384 to Val, known to be operable in yeast ALS and Trp 591 to Leu, known to be operable in yeast ALS and ALS encoded by the SURB-Hra gene of tobacco. By combining the above mutations, double mutations, resulting in two amino acid substitutions such as Ala 122 to Ser and Pro 197 to Ser, or Ala 122 to Ser and Pro 197 to Ala, or Pro 197 to Ala and Trp 591 to Leu, or Pro 197 to Ser and Trp 591 to Leu have also been made. These mutations were made in a DNA fragment that included only about 180 bp in the 5' direction (upstream) and only about 600 bp in the 3' direction (downstream) of the ALS coding sequence. These DNA fragments were introduced into tobacco by transformation. Herbicide resistance was not expressed in these transformants. It is believed that utilization of the regulatory sequences found sufficient for expression of herbicide resistance with the SURA-C3 mutant coding region, namely 2.5 kb in the 5' direction (upsteam) and 1.8 kb in the 3' direction (downstream) of the coding region, would have resulted in expression of herbicide resistance with all of the site-directed mutations in the SURA coding region.

Site directed mutations that are expected to code for herbicide resistant ALS have also been made in the sugarbeet ALS gene. These mutations result in the following amino acid substitutions: Ala 122 to Pro, known to be operable in yeast ALS, Pro 197 to Ala, known to be operable in ALS encoded by the SURB-Hra gene of tobacco, Trp 591 to Leu, known to be operable in yeast ALS and in ALS encoded by the SURB-Hra gene of tobacco and the double mutant, Pro 197 to Ala and Trp 591 to Leu, known to be operable in ALS encoded by the SURB-Hra gene of tobacco.

The nucleic acid fragments of the invention generally can be introduced into plants directly or in a nucleic acid construct comprising the desired nucleic acid fragment. The nucleic acid construct can be derived from a bacterial plasmid or phage, from the Ti- or Ri-plasmids, from a plant virus or from an autonomously replicating sequence. One preferred means of introducing the nucleic acid fragment into plant cells comprises use of Agrobacterium tumefaciens containing the nucleic acid fragment between T-DNA borders either on a disarmed Ti-plasmid (that is, a Ti-plasmid from which the genes for tumorigenicity have been deleted) or in a binary vector in trans to a Ti-plasmid with Vir functions. The Agrobacterium can be used to transform plants by inoculation of tissue explants, such as stems or leaf discs, or by co-cultivation with plant protoplasts. Another preferred means of introducing the present nucleic acid fragment comprises direct introduction of the fragment or a vector containing the fragment into plant protoplasts or cells, with or without the ald of electroporation, polyethylene glycol or other agents or processes known to alter membrane permeability to macromolecules.

The nucleic acid fragments of the invention can be used to transform protoplasts or cell cultures from a wide range of higher plant species to form plant tissue cultures of the present invention. These species include the dicotyledonous plants tobacco, petunia, cotton, sugarbeet, potato, tomato, lettuce, melon, sunflower, soybean, canola (rapeseed) and other Brassica species and poplars; and the monocotyledonous plants corn, wheat, rice, Lolium multiflorum and Asparagus officinalis. It is expected that all protoplast-derived plant cell lines can be stably transformed with the fragments of the invention can also be introduced into plant cells with subsequent formation of transformed plants of the present invention. Transformation plants of the present invention. Transformation of whole plants is accomplished in plants whose cells can be transformed by foreign

genes at a stage from which whole plants can be regenerated. In the present invention, transformed plants are monocotyledonous and dicotyledonous plants. Preferably, the transformed plants are selected from the group consisting of tobacco, petunia, cotton, sugarbeets, potato, tomato, lettuce, sunflower, soybean, canola and other Brassica species, poplars, alfalfa, clover, sugarcane, barley, oats and millets; see "Handbook of Plant Cell Culture" Vols. 1-3, Evans, D. A. et al., Sharp et al., and Ammirato et al., respectively, MacMillan, N.Y. (1983, 84). Most preferably, the transformed plants are selected from the group consisting of tobacco, petunia, potato, tomato, sunflower, sugarbeet, alfalfa, lettuce or Brassica species. The range of crop species in which foreign genes can be introduced is expected to increase rapidly as tissue culture and transformation methods improve and as selectable markers such as the fragments of the invention (see discussion below) become

One could further increase the level of expression of the nucleic acid fragments of the invention by replacing their native regulatory nucleotide sequences, 5' and 3' to the ALS coding sequence, with synthetic or natural sequences known to provide high level and/or tissue specific expression. One may also substitute the nucleotide sequences of the nucleic acid fragments of the invention with other synthetic or natural sequences which encode transit peptides which will allow efficient chloroplast uptake of the nucleic acid fragments of the invention

The nucleic acid fragments of the present invention also have utility as selectable markers for both plant genetic studies and plant cell transformations. A gene of interest, generally conferring some agronomically useful trait, e.g. disease resistance, can be introduced into a population of sensitive plant cells physically linked to a nucleic acid fragment of the present invention. Cells can then be grown in a medium containing a herbicide to which the ALS encoded by a fragment of the invention is resistant. The surviving (transformed) cells are presumed to have acquired not only the herbicide resistance phenotype, but also the phenotype conferred by the gene of interest. The nucleic acid fragments can be introduced by cloning vehicles, such as phages and plasmids, plant viruses, and be direct nucleic acid introduction. Subsequently, in a plant breeding program, the agronomically useful trait can be introduced into various cultivars through standard genetic crosses, by following the easily assayed herbicide resistance phenotype associated with the linked selectable genetic marker.

Transformed plants of the present invention are resistant to many of the sulfonylurea, triazolo pyrimidine sulfonamide and imidazolinone herbicides. These herbicides are disclosed in the following patents and published patent applications as follows:

published patent applications as follows.	
Sulfonylureas U.S. 4,127,405 U.S. 4,383,113	
U.S. 4,169,719 U.S. 4,394,153	
U.S. 4,190,432 U.S. 4,394,506	
U.S. 4,214,890 U.S. 4,420,325	
U.S. 4,225,337 U.S. 4,452,628	
U.S. 4,231,784 U.S. 4,481,029	
11.5 4.257.802 11.5 4.586.950	

U.S. 4,310,346 U.S. 4,514,401 U.S. 4,435,206 U.S. 4,634,465 U.S. 4,634,465

EP-A-204,513

<u>Triazolopyrimidine sulfonamides</u>
South African Application 84/8844 (published 5/14/85)

Imidazolinones

U.S. 4,188,487 EP-A-41,623 (published 12/16/81)

The nucleic acid fragments of the present invention encode ALS which is resistant to the following sulfonylurea herbicides:

I

wherein R is H or CH3:

65

10

25

30

40

45

50

55

J is 
$$R_2$$
 $R_2$ 
 $R_1$ 
 $R_2$ 
 $R_1$ 
 $R_2$ 
 $R_1$ 
 $R_2$ 
 $R_1$ 
 $R_2$ 
 $R_3$ 
 $R_4$ 
 $R_4$ 
 $R_5$ 
 $R_7$ 
 $R_6$ 
 $R_6$ 

$$R_8$$
  $P^n$ :  $R_8$   $P^n$  or  $R_{20}$   $P^n$ :  $R_{20}$   $P^n$ :  $P^n$ 

 $R_1$  is Cl, Br, NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>2</sub>-C<sub>4</sub> alkenyl, CF<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkoxy, C<sub>3</sub>-C<sub>4</sub> alkynyloxy, CO<sub>2</sub>R<sub>8</sub>, CONR<sub>10</sub>R<sub>11</sub>, S(O)<sub>m</sub>R<sub>12</sub>, OSO<sub>2</sub>R<sub>12</sub>, phenyl, SO<sub>2</sub>N(OCH<sub>3</sub>)CH<sub>3</sub>, SO<sub>2</sub>NR<sub>10</sub>R<sub>11</sub>,

30

35

40

R2 is H, Ci, Br, F, CH<sub>3</sub>, NO<sub>2</sub>, SCH<sub>3</sub>, OCF<sub>2</sub>H, OCH<sub>2</sub>CF<sub>3</sub> or OCH<sub>3</sub>; 60  $R_3 \, \text{is Cl, NO}_2, CO_2 CH_3, CO_2 C_2 H_5, SO_2 N (CH_3)_2, SO_2 CH_3 \, \text{or SO}_2 C_2 H_5; \\$  $R_4 \text{ is } C_1 - C_3 \text{ alkyl}, Cl, Br, NO_2, CO_2 R_9, CON(CH_3)_2, SO_2N(CH_3)_2, SO_2N(OCH_3)CH_3 \text{ or } S(O)_m R_{12}; \\$  $R_5\,is\,C_1-C_3\,alkyl,\,C_4-C_5\,cycloalkylcarbonyl,\,F,\,Cl,\,Br,\,NO_2,\,CO_2R_{14},\,SO_2N(CH_3)_2,\,SO_2R_{12}\,or\,phenyl;$ R8 is H, C1-C3 alkyl, or CH2CH = CH2; R7 is H, CH3, OCH3, Cl or Br; 65

$R_0$ is $C_1$ - $C_4$ alkyl, $C_3$ - $C_4$ alkenyl or $C_1$	
$R_{10}$ is H or $C_1$ - $C_3$ alkyl;	
R <sub>11</sub> is H or C <sub>1</sub> -C <sub>2</sub> alkyl;	
R <sub>12</sub> is C <sub>1</sub> -C <sub>3</sub> alkyl:	
R <sub>13</sub> is H or CH <sub>3</sub> ;	5
$R_{14}$ is $C_1$ - $C_3$ alkyl or $CH_2CH = CH_2$ ;	
mis 0, 1 or 2;	
nis1or2;	
Q is CH <sub>2</sub> , CHCH <sub>3</sub> or NR <sub>15</sub> ;	
R <sub>15</sub> is H or C <sub>1</sub> -C <sub>4</sub> alkyl;	10
P is O or CH₂;	
Rasis Hor CHa;	
H <sub>17</sub> is C(O)NR <sub>18</sub> R <sub>18</sub> :	
R <sub>18</sub> is H or CH <sub>3</sub> ;	
	15
R <sub>19</sub> is CH <sub>3</sub> ;	
R <sub>20</sub> is H, Cl, F, Br, CH <sub>3</sub> , CF <sub>3</sub> , OCH <sub>2</sub> or OCF <sub>2</sub> H;	
R <sub>21</sub> is H or CH <sub>3</sub> ;	
X is CH <sub>3</sub> , OCH <sub>3</sub> , OC <sub>2</sub> H <sub>5</sub> or NHCH <sub>3</sub> ;	
Y is CH <sub>3</sub> , C <sub>2</sub> H <sub>5</sub> , OCH <sub>3</sub> , OC <sub>2</sub> H <sub>5</sub> , OCF <sub>2</sub> H, OCH <sub>2</sub> CF <sub>3</sub> , Cl, CH <sub>2</sub> OCH <sub>3</sub> or cyclopropyl;	
Z is CH or N;	20
and their agriculturally suitable salts; provided that	
a) when Y is CI, then Z is CH and X is OCH₃;	
b) when Y is OCF <sub>2</sub> H, then Z is CH;	
c) when J is J-1 and R <sub>1</sub> is OSO <sub>2</sub> R <sub>12</sub> or phenyl, then Y is other than OCF <sub>2</sub> H;	
d) when J is J-2, then Y is other than OCF <sub>2</sub> H or OCH <sub>2</sub> CF <sub>3</sub> ; and	25
g) when J is J-2, then Y is other than Oct 2110 oct 120 3, and	
e) when J is J-3 and R <sub>4</sub> is S(O) <sub>m</sub> R <sub>12</sub> , then Y is other than OCH <sub>2</sub> CF <sub>3</sub> .	
Sulfonylurea herbicides to which the ALS is particularly resistant include	
1) Compounds of Formula I where J is J-1;	
R <sub>1</sub> is Cl, CH <sub>3</sub> , C <sub>1</sub> -C <sub>4</sub> alkoxy, C <sub>1</sub> -C <sub>2</sub> haloalkoxy, allyloxy, propargyloxy, CO <sub>2</sub> R <sub>8</sub> , CONR <sub>10</sub> R <sub>11</sub> ,	
SO <sub>2</sub> N(OCH <sub>3</sub> )CH <sub>3</sub> , SO <sub>2</sub> NR <sub>10</sub> R <sub>11</sub> , S(O) <sub>m</sub> R <sub>12</sub> , OSO <sub>2</sub> R <sub>12</sub> , phenyl or	30
n h CH3	35
2) Compounds of Formula I where	40
Jis J-2;	
R is H; and	
R3 IS SO2N(CH3)2, CO2CH3 or CO2C2H5.	
3) Compounds of Formula I where	
JisJ-3	
Ris H; and	45
R <sub>4</sub> is CO <sub>2</sub> CH <sub>3</sub> or CO <sub>2</sub> C <sub>2</sub> H <sub>5</sub> ;	
4) Compounds of Formula I where	
JisJ-4;	
RisH;	50
R <sub>5</sub> is CI, Br, CO <sub>2</sub> CH <sub>3</sub> , CO <sub>2</sub> C <sub>2</sub> H <sub>5</sub> or	
0	
n	
-c-<   :	
7	55
R <sub>6</sub> is CH <sub>3</sub> ; and	
R7 is H, Clor OCH₃;	
5) Compounds of Formula I where	
J is J-5:	60
RisH;	
R <sub>5</sub> is CO <sub>2</sub> CH <sub>3</sub> or CO <sub>2</sub> C <sub>2</sub> H <sub>5</sub> ; and	
R <sub>7</sub> is H or CH <sub>3</sub> .	•
6) Compounds of Formula I wherein	6
Jis J-6:	

```
Q is CHCH3 or NR15;
                                            R is H; and
                                            Rs is H, F, Cl, CH<sub>3</sub>, OCH<sub>3</sub>, CF<sub>3</sub> or SCH<sub>3</sub>.
                                                       7) Compounds of Formula I where
                                              J is J-7;
 5
                                             R is H;
                                              P is O; and
                                              R<sub>8</sub> is H, F, Cl, CH<sub>3</sub>, OCH<sub>3</sub>, CF<sub>3</sub> or SCH<sub>3</sub>.
                                                         8) Compounds of Formula I where
                                              Jis J-8;
10
                                               R is H:
                                               R<sub>18</sub> is CH<sub>3</sub>; and
                                               R<sub>8</sub> is H, F, Cl, CH<sub>3</sub>, OCH<sub>3</sub>, CF<sub>3</sub> or SCH<sub>3</sub>.
                                                        9) Compounds of Formula I where
                                                J is J-9;
15
                                               R is H; and
                                                R<sub>17</sub> is C(O)N(CH<sub>3</sub>)<sub>2</sub>.
                                                            10) Compounds of Formula I where R is H;
                                                R_1 \text{ is Cl, C}_1\text{-C}_4 \text{ alkoxy, OCF}_2\text{H, OCH}_2\text{Cl, CO}_2\text{R}_9, \text{CON(CH}_3)_2, \text{SO}_2\text{N(CH}_3)_2, \text{SO}_2\text{R}_{12} \text{ or OSO}_2\text{R}_{12}; \text{ and } \text{CON(CH}_3)_2, \text{CON(CH}_3)_
                                                 R2 is H, C1, CH3, or OCH3.
20
                                      The nucleic acid fragments of the present invention encode ALS which is resistant to the following
                            triazolopyrimidine sulfonamides:
 25
   30
                                                                                                     II
    35
                               wherein
                                Ar is
      45
                                   Re is C1-C4 alkyl, F. Cl. Br. I, NO2, S(O)pRd, COORe or CF3;
```

Rb is H, F, Cl, Br, I, C1-C4 alkyl or COORe;

Re is H, C1-C4 alkyl, F, Cl, Br, I, CH2ORd, phenyl, NO2 or COORe;

Rd is C1-C4 alkyl;

Re is C1-C4 alkyl, C1-C4 alkenyl, C1-C4 alkynyl, or 2-ethoxyethyl;

V is H, C1-C3 alkyl, allyl, propargyl, benzyl or C1-C3 alkylcarbonyl;

X1, Y1, and Z1, are independently H, F, Cl, Br, I, C1-C4 alkyl C1-C2 alkylthio or C1-C4 alkoxy; and

D

Triazolopyrimidinesulfonamide herbicides to which the ALS is particularly resistant include

1) Compounds of Formula II where V is H.

2) Compounds of Preferred 1 where

X<sub>1</sub> is H or CH<sub>3</sub>;

Y1 is H;

 $Z_1$  is  $\text{CH}_3$  ; and  $R_e$  and  $R_c$  are not simultaneously H.

The nucleic acid fragments of the present invention encode ALS which is resistant to the following imidazolinones:

65

*55* ,

wherein

III

M O N N

R<sub>f</sub> is C<sub>1</sub>-C<sub>4</sub> alkyl;

Rg is C1-C4 alkyl or C3-C6 cycloalkyl;

A1 is COORI, CH2OH or CHO;

30

10

15

20

25

R<sub>I</sub> is H; C<sub>1</sub>-C<sub>12</sub> alkyl optionally substituted by C<sub>1</sub>-C<sub>3</sub> alkyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl or phenyl; C<sub>3</sub>-C<sub>6</sub> alkenyl optionally substituted by phenyl or 1-2 C<sub>1</sub>-C<sub>3</sub> alkyl, F, Cl, Br, or I; or C<sub>3</sub>-C<sub>5</sub> alkynyl optionally substituted by phenyl or 1-2 C<sub>1</sub>-C<sub>3</sub> alkyl, F, Cl, Br or I;

B is H; C(O)C1-C6 alkyl or C(O)phenyl optionally substituted by Cl, NO2 or OCH3;

X2 is H. F. Cl. Br. I. OH or CH3;

35

Y<sub>2</sub> and Z<sub>2</sub> are independently H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, F, Cl, Br, I, phenyl, NO<sub>2</sub>, CN, CF<sub>3</sub> or SO<sub>2</sub>CH<sub>3</sub>; X<sub>3</sub> is H, C<sub>1</sub>-C<sub>3</sub> alkyl, F, Cl, Br, I or NO<sub>2</sub>; and

L, M, Q and Rh are independently H, F, Ci, Br, I, CH<sub>3</sub>, OCH<sub>3</sub>, NO<sub>2</sub>, CF<sub>3</sub>, CN, N(CH<sub>3</sub>)<sub>2</sub>, NH<sub>2</sub>, SCH<sub>3</sub> or SO<sub>2</sub>CH<sub>3</sub> provided that only one of M or Q may be a substituent other than H, F, Cl, Br, I, CH<sub>3</sub> or OCH<sub>3</sub>.

Imidazolinone herbicides to which the ALS is particularly resistant include

40

45

· 50

60

65

1) Compounds of Formula III where

B is H; and

A1 is COORI.

2) Compounds of Preferred 1 where

Ris CH3;

Rg is CH(CH<sub>3</sub>)<sub>2</sub>;

X2 is H;

Y2 is H, C1-C3 alkyl or OCH3;

Z2 is H:

X<sub>3</sub> is H, CH<sub>3</sub>, Cl or NO<sub>2</sub>; and

L, M, Q and Rh are H.

Any of the aforementioned compounds may be applied alone or in combination to the site, pre-and/or post-emergence. Because the crop plant itself is resistant to the herbicide(s), the spectrum of herbicide activity can be chosen for its efficacy in controlling the unwanted vegetation.

The present invention is further defined in the following Examples, in which all parts and percentages are by weight and degrees are Celsius; unless otherwise stated. It should be understood that these examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above disclosure and these examples one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

**EXAMPLE I** 

Tobacco (Nicotiana tabacum cv. Xanthi) DNA from the Hra mutant was made according to the procedure of Dunsmuir et al. (J. Mol. App. Genetics, 1983, 2, 285). 2 x 13 g of 1-2 inch tobacco leaves were removed from plants and immediately ground in 2 x 20 mL buffer A (10 mM Tricine-KOH pH 7.6 - 1.14 M sucrose - 5 mM

MgCl2 - 5 mM 2-mercaptoethanol) in the cold room, using mortars and pesties. An additional 40-50 mL of buffer A was added, and the slurries were filtered through 16 layers of cheesecloth. The filtrates were centrifuged at 2500 rpm in a Sorvall GSA rotor at 4°C for 5 minutes. The pellets were resuspended in 10 mL buffer A, another 100 mL of buffer A was mixed in, and the cells were centrifuged as above. The pellets were then resuspended in 100 mL buffer A + 0.4% Triton X-100, and left on ice for 10 minutes; and centrifuged as above. The pellets were washed twice more in the latter buffer. The final pellets were resuspended in 5 mL of resuspension buffer (50 mM Tris HCl pH 8, 20 mM EDTA), 1 mL of resuspension buffer, 10% sarkosyl was added, and the volumes were then adjusted to 10 mL with suspension buffer. Proteinase K was added to 100 μg/mL to the lysates, and the lysates were digested at 37°C overnight. The lysates were then brought to a density of 1.55 g/mL CsCl, and to a final concentration of 300 µg/mL ethidium bromide. The solutions were centifuged in a Beckman Ti70.1 rotor at 40000 rpm at 15°C for 24 hours, and the fluorescent DNA band was removed after visualization with longwave UV light. To remove the DNA, holes were punched in the sides of the polyallomer tubes with an 18 gauge needle, and the viscous DNA was allowed to drip into collection tube. Great care was taken at all stages after cell lysis to prevent shearing of the DNA. The DNA was again gently resuspended in a CsCl solution of 1.55 g/mL density and 300 µg/mL ethidium bromide, and centrifuged at 40000 rpm at 15°C for 48 hours, in a Sorvall TFT65.13 rotor. The DNA was again collected by side puncture of the tube. It was gently extracted 10 times with TE (10 mM Tris HCl pH 8, 1 mM EDTA) saturated-isoamyl alcohol, and then dialyzed extensively against TE.

The standard techniques of recombinant DNA and molecular cloning used here are described in R. W. Davis, D. Botstein and J. R. Roth, Advanced Bacterial Genetics, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1980) and T. Maniatis, E. F. Fritsch and Sambrook, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1982).

A tobacco DNA library was constructed following the procedures of Maniatis et al (see above). Tobacco DNA was digested with the restriction enzyme Sau 3A to give a majority of fragments in the 20 Kilobase size range, as assayed by agarose gel electrophoresis. The fragments were loaded onto 10-40% sucrose (in 1 M NaCl, 20 mM Tris pH 8, 1 mM EDTA) gradients and size-fractionated by centrifugation in a Beckman SW28 rotor at 26000 rpm at 17°C for 16 hours. Fractions from the sucrose gradients were collected and analyzed by agarose gel electrophoresis, and fractions containing fragments in the 20 kilobase size range were dialyzed against TE and ethanol precipitated. They were then ligated to BamH I cut phage lambda EMBL3 arms, at a 2:1 molar ratio, and packaged into lambda phage heads, following the instructions supplied by the manufacturer of the lambda arms and packaging reactions (Stratagene Cloning Systems, San Diego, CA).

A tobacco DNA library of 400000 phage was plated on the host strain E. coli LE 392 (Silhavy, T. J., Berman, M. L. and Enquist, L. W. (1984), "Experiments with Gene Fusions," Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.) at a density of 50000 phage per 150 mm Petri dish, on 10 plates. Duplicate nitrocellulose filter lifts of the phage plaques were made according to the procedure of Maniatis et al., and were hybridized with \$^2P\$-labeled probes carrying either 5' or 3' ALS gene fragments produced in a riboprobe labeling system. Riboprobes were synthesized according to the procedures accompanying the riboprobe kit sold by Promega Biotech (Madison, WI). Plaques that gave positive signals on films from both sets of filters were picked and the purification process was reiterated, until well-isolated hybridizing plaques were obtained.

Minipreps of the DNA from plaque purified phage were analyzed by restriction enzyme digestions. Two classes of cloned tobacco DNA fragment inserts were distinguished as shown in Figure 1. Phages 1, 2, 17 and 18 contained inserts related to the previously isolated ALS gene from the SURA locus, encoding herbicide sensitive ALS. Phage 3 contained an insert distinct from the above which was expected to contain the SURB-Hra gene encoding herbicide resistant ALS. EcoR I fragments that encompassed the hybridizing regions of phage 3 were subcloned into M13 phage vectors and subjected to DNA sequence analysis, using oligonucleotides to extend the sequenced regions in overlapping segments. A single open reading frame of 1992 nucleotides was found, and was identified as an ALS gene by comparison of the deduced amino acid sequence with conserved regions of the amino acid sequences of ALS proteins from other species.

ALS genes Isolated from the herbicide-resistant mutant tobacco, Hra, were introduced into sensitive tobacco cells via the "binary vector" system employing Agrobacterium tumefaciens. The ALS genes were first introduced into a binary vector in A. tumefaciens via plasmid conjugation, and the engineered A. tumefaciens were then used to transform plant cells with the foreign genes via co-cultivation.

A) Introduction of the Isolated Tobacco ALS Genes into A. tumefaciens:

ഞ

65

i) Construction of Binary Vectors: The standard techniques of recombinant DNA and molecular cloning used here are described in R. W. Davis, D. Botstein and J. R. Roth, Advanced Bacterial Genetics, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1980) and T. Maniatis, E. F. Fritsch and Sambrook, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1982). The purified 8.3 kilobase Spe I nucleic acid fragment of the Invention, which was isolated from the Hra tobacco mutant and which contains a coding sequence for a herbicide-resistant form of an ALS gene, was inserted into the Xba I site of the plasmid vector pMuc19 (J. D. G. Jones, P. Dunsmiur and J. Bedbrook, EMBO Journal 4:2411-2418 (1985)). (Although, Spe I and Xba I restriction enzymes recognize different DNA sequences, the products of these digestions carry the same 5' overhanging sequence). The orientation of the insert fragment in one of the resultant plasmids, pAGS148, was determined by restriction enzyme analyses (Fig. 2).

The binary vector pAGS135 was used to move plasmid pAGS148 into A. tumefaciens. Plasmid pAGS135 is

derived from plasmid pAGS112 (P. Van den Elzen, K. Y. Lee, J. Townsend and J. Bedbrook, Plant Mol. Biol., 5:149-154 (1985)) by digestion of plasmid pAGS112 DNA with Xho I restriction endonuclease, treatment with the Klenow fragment of E. coll DNA polymerase I, and self-ligation of the DNA which effects the removal of the Xho I site outside the T-DNA right border. Plasmid pAGS112 is derived fro the wide-host range vector pLAFR (A. M. Friedan, S. R. Long, S. E. Brown, S. E. Buikema and F. M. Ausubel, Gene, 18:289-296 (1982)) by the insertion into pLAFR of an EcoR I fragment in which the T-DNA borders flank a gene for expressing kanamycin resistance in plants and a unique BamH I site for cloning (Van den Elzen et. al., Plant Mol. Biol., 5:149-154 (1985)). CsCl purified plasmids pAGS148 and pAGS135 were digested with BamH I, and the resultant BamH I-cleaved plasmids were ligated. The ligation mixtures were packaged into lambda phage particles in vitro and used to infect Escherichia coli strain HB101. Transformants were selected on amplcillin. The physical map of a recombinant plasmid, pAGS152, from one of the transformants was determined by restriction analyses and is shown in Fig. 2.

10

15

20

25

30

35

40

45

50

55

ii) Conjugation of Plasmid pAGS152 from E. coli into A. tumefaciens: Plasmid pAGS152 was introduced into A. tumefaciens by conjugation essentially by the three-way mating method of Ruvkun, G. and Ausubel, F. M., Nature, 289:85–88 (1981). E. coli strain HB101 harboring plasmid pAGS152 and E. coli strain HB101 harboring the mobilizing vector pRK2013 (ATCC 37159) (D. Figurski and D. R. Helinski, Proc. Natl. Acad. Sci U.S.A., 76:1648-1652 (1979)) were mixed with A. tumefaciens strain LBA4404 harboring plasmid pAL4404 (A. Hoekema, P. R. Hirsch, P. J. J. Hooykaas and R. A. Schilperoort, Nature, 303:179-180 (1983)) and allowed to mate on solid LB medium (J. H. Miller, Experiments in Molecular Genetics, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1972)) at 28°C for 16 hours. Transconjugants were selected on plates containing rifampicin at 100 mg/liter and tetracycline at 1 mg/liter. A. tumefaciens LBA4404:pAGS152 was restreaked on minimal A medium containing tetracycline at 1 mg/liter.

Essentially, a similar method was used to obtain both A. tumefaciens LBA4404 containing plasmid pAGS112, the binary vector without any plant nucleic acid fragment insert, and A. tumefaciens LBA4404 containing plasmid pAGS145, the binary vector containing a nucleic acid fragment from phage clone 1. The latter fragment was also isolated from Hra mutant tobacco plants and carries a gene for a herbicide-sensitive form of ALS; this gene is not the wild type allele of the gene for the herbicide-resistant ALS in the nucleic acid fragment of the invention but the SURA gene from the second genetic locus.

B) Introduction of the Isolated ALS Genes into Sensitive Tobacco by Co-cultivation of the Plant Cells with A. tumefaciens LBA4404 (pAGS145) and LBA4404 (pAGS152).

All manipulations of sterile media and plant materials were done in laminar flow hoods, under suitable containment. Plant growth and plant cell cultures were carried out at 27°C. All protoplast manipulations were carried out at room temperature unless otherwise mentioned.

Day 1 (afternoon):

Protoplast isolation medium was prepared by adding the following to K3/S(1) Medium: 0.1% (w/v) of MES buffer (Sigma Chemical Co.), 1% (w/v) of Cellulase (Onozuka or Cellulysin), and 0.1% of Macerase (Onozuka). After gentle stirring for approximately 30 minutes, the pH was brought to 5.6 with KOH and the medium was filter-sterized.

Sterile tobacco (Nicotiana tabacum var. Wisconsin 38) plants were cultured from 1 cm apical or auxillary explants on OMS Medium in Magenta Boxes under a cycle of a 16 hour light period (6,000-8,000 lux) followed by an 8 hour dark period. When the plants were 5-7 weeks old, fully expanded leaves (3-6 leaves down from the apex) were removed, and two leaves each were placed, top surface down, on 20 mL of protoplast isolation medium in a 100 x 25 mm petri dish. The leaves were then submerged and finely divided with a sharp surgical blade. The midrib was held and the cuts were made outward from it towards the leaf margin at approximately 2 mm intervals. The petri dishes were then sealed with parafilm and the macerated tissue incubated overnight (14-17 hours) in darkness at 27-29°C with gentle gyrotory agitation (20-25 rpm).

Day 2 (moming):

A 75 mm filtering funnel, lined with four layers of cheesecloth, was clamped to a ringstand. A glass tube (approximately 15 cm long and with an outer diameter of < 5 mm) was attached to the funnel with latex tubing. The funnel, cheesecloth, latex and glass tubing were wrapped in aluminum foil and sterilized in an autoclave as

The glass tubing was placed in a Babcock bottle, and the cheesecloth was wetted with K3/S(1) Medium. The digested leaf tissue from two petri dishes was carefully poured into the funnel. The cheesecloth was rinsed and expressed into the bottle. The loaded bottles were topped with K3/S(1) Medium, covered with foil and centrifuged at approximately 100 x g for 10 minutes. The floating protoplasts (1-2 mL) were collected with a 1 mL serological pipette and placed in 20 mL of K3/S(1) Medium in another Babcock bottle. After resuspending the protoplasts by gently swirling, the Babcock bottles were topped and centrifuged as before. The floating protoplasts (1-2 mL) were collected as described above and placed in 30 mL of K3/G(1) Medium. The protoplasts were counted in a hemacytometer, and the volume was adjusted to give 1 x 10+5 protoplasts/mL. 5 mL aliquots of the protoplasts were plated in petri dishes [100 x 20 mm tissue-culture petri dishes (Corning); these dishes were used in all subsequent protoplast manipulations] and cultured in darkness.

Day 2 (afternoon):

A single colony of A. tumefaciens, containing and desired plant transformation vector, viz., pAGS112 (plasmid vector above), pAGS152 (containing the nucleic acid fragment of the present invention) or pAGS145 (containing a nucleic acid encoding a sensitive form of ALS), growing on a Minimal A plate was inoculated into 5 mL of Minimal A Medium in an 18 mm test tube and cultured overnight on a roller drum at 40-60 rpm at 27-28°C.

Day 3 (morning):

10

20

The optical density of the A. tumefaciens cultures was measured at 550 nm and adjusted to 0.15 with Minimal A Medium, and the bacteria were allowed to continue growing as described above.

Day 3 (afternoon):

When the optical density (at 550 nm) of the A. <u>tumefaciens</u> cultures was 0.6 (log phase culture), approximately 6 hours after dilution, the bacteria were added to plant cells at a multiplicity of approximately 50 bacteria/plant cell (an optical density of 1.0 at 550 nm = 1.4 x 10<sup>9</sup> bacteria). The bacteria and plant cell mixture was co-cultivated for 66 hours at 24°C in low light (approximately 500 lux). Non-transformed protoplast controls were incubated similarly, but without agrobacteria. The following protocol is carried out for each co-cultivation (transformed cells with different agrobacteria, as well as non-transformed cells).

Day 6 (morning):

Co-cultivation was terminated by adding 20 mL of a 1:1 mixture of K3/G(2) Medium:C Medium supplemented with 500 mg/liter of cefotaxime (to select against the agrobacteria) to 5 mL of the co-cultivation mixture. The co-cultivated cells were gently and thoroughly resuspended in the new medium by mixing with a 5 or 10 mL serological pipette. The cell density was 2 x 10<sup>4</sup> protoplast equivalents/mL (protoplast equivalents = Initial protoplasts, assuming 100% recovery and cell survival) and the osmoticum was 0.35 M. Three 5 mL aliquots of each culture were dispensed into fresh petri dishes.

From this juncture until the cells were embedded in solid medium, the cells were cultured in low light (500-1500 lux) without motion and were aseptically transferred to different media. At the indicated times, cells from one plate of each culture were transferred to non-selective media, while cells from the other two plates of each culture were transferred to selective media containing either 50 mg/liter of kanamycin or 2 ng/mL chlorsuffuron in order to select for transformed plant cells. For these transfers, the contents of each plate were collected with a 5 mL serological pipette, placed in separate 15 mL polystyrene conical centrifuge tubes and centrifuged at approximately 50 x g for 5-10 minutes. The supernatant fluid was removed with a pipette without disrupting the loose pellet. Pellets of co-cultivated cells from each plate were then gently resuspended in the appropriate fresh medium.

Day 10:

The cells were transferred into 5 mL of C Medium supplemented with 500 mg/liter cefotaxime in the case of non-selected plant cells or with 500 mg/liter cefotaxime and either 50 mg/liter of kanamycin or 2 ng/mL chlorsulfuron in the case of selected cells. Each of these cultures was returned to the petri dishes from which they were taken; in this way not all cells needed to be pelleted to effect a medium exchange with minimal cell loss.

Day 13:

Non-selected cells were transferred to 20 mL of a 3:1 mixture of C Medium:MSP Medium supplemented with 500 mg/liter of cefotaxime, and a 5 mL aliquot was dispensed into a fresh petri dish (at a density of 5 x 10<sup>3</sup> protoplast equivalents/mL). The selected cells were resuspended in 5 mL of a 3:1 mixture of C Medium:MSP Medium supplemented with 500 mg/liter cefotaxime and 50 mg/liter kanamycin or 2 ng/mL chlorsulfuron and returned to the original plates for further culture.

ŝ

Day 16-17:

The cells were transferred to 5 mL of a 1:1 mixture of C Medium and MSP Medium supplemented with 500 mg/liter cefotaxime alone (In the case of non-selected plant cells) or with 500 mg/liter cefotaxime and either 50 mg/liter of kanamycin or 2 ng/mL chlorsulfuron (in the case of selected cells) and cultured as before.

Day 20:

55

The non-selected cells were transferred to 25 mL of 1:1 mixture of C Medium:MSP Medium, and the mixture added to 25 mL of a 1:1 mixture of a 2 X MSP Medium and 1% (w/v) type VII agarose solution (50°C). The resultant culture was mixed quickly with a 25 mL wide-mouth serological pipette and dispensed in 5 mL aliquots into fresh petri dishes. The suspended micro calluses in the agar solution were spread carefully and evenly across the plates with agitation by hand. The plates were covered and left in the hood for one hour to solidity before they were wrapped in parafilm and removed to the culture chamber. The cell density was about 5 X 10² protoplasts equivalents/mL and the osmoticum was 0.15 M. The embedded cells were counted on a colony counter approximately 10 days later (Tables 1 and 2, below).

The selected cells were transferred to 20 mL of a 1:3 mlxture of C Medium:MSP Medium containing 50 mg/liter of kanamycin or 2 ng/mL chlorsulfuron. Five mL aliquots of the resuspended cultures (5 X  $10^{+3}$  protoplast equivalents/mL) were dispensed into four fresh petrl dishes per selected culture and cultured as before.

Day 23-24:

Each 5 mL culture of the selected cells was diluted with 7.5 mL of MSP Medium supplemented with 50 mg/liter of kanamycin or 2 ng/mL chlorsulfuron in order to achieve a cell density of 2 X 10+3 protoplast equivalents/mL. This density-adjusted culture was mixed with 12.5 mL of a 1:1 mixture of 2X MSP Medium and 1.0% (w/v) type VII agarose solution (50°C) supplemented with 50 mg/liter of kanamycin or 2 ng/mL chlorsulfuron. Five mL aliquots of the mixed cultures were quickly dispensed with a 25 mL wide-mouth serological pipette into fresh petri dishes. The final plating density was 1 X 10³ protoplast equivalents/mL, and the osmoticum of the culture was 0.1 M. The plates were solidified as described above. The embedded cells were scored for growth on a colony counter approximately 10 days later.

Day 25+:

Ten to twelve individual transformed calluses/colonies were picked and transferred with a No. 11 scalpel to a petril dish containing MSR medium with or without the appropriate selective agent for plant regeneration. The calluses were cultured at 27°C, with a photo period of 16 hours of light (5000-8000 lux) followed by 8 hours of darkness. Shoots appeared after 2-3 weeks and continued to be produced for several months. Shoots of 2-3 mm length were excited with a sharp surgical blade and transferred for rooting to OMS Medium in Magenta houses.

After root formation (1 to 4 weeks), plants were transferred to soil for regeneration by the methods of R. S. Chaleff and M. F. Parsons, Proc. Natl. Acad. Sci.-U.S.A. 75:5104 (1978), and B. Tisserat in Plant Cell Culture: A Practical Approach, Ed. Dixon, R. A., IRL Press, Oxford (1985).

Results of Co-cultivation:

The results of the co-cultivation experiments show that the nucleic acid fragment of the invention — but not the tobacco SURA gene for the herbicide-sensitive form of ALS — confers herbicide resistance when introduced into herbicide-sensitive tobacco cells (Tables 4 and 5, below). Since the nucleic acid fragment of the invention can confer herbicide resistance at a similar frequency when introduced in either orientation with respect to the vector, it is believed to contain the regulatory sequences both 5' and 3' to the coding sequence which are required for the expression of the herbicide-resistant ALS gene.

The level of herbicide resistance conferred by the nucleic acid fragment of the invention was determined by plating one hundred colonies each of pAGS152 transformed N. tabacum cells resistant to chlorsulfuron at 2 ppb and non-co-cultured wild type N. tabacum cells on different concentrations of chlorsulfuron. The number of colonies actively growing on different concentrations of chlorsulfuron after one month was scored (Table 6, below). While wild type colonies are sensitive to chlorsulfuron at 2 ppb, colonies derived from co-cultivation with A. tumefaciens containing pAGS152 could tolerate up to 2000 ppb. This level of resistance of the transformants is comparable to that of the Hra herbicide-resistant mutant tobacco from which the nucleic acid fragment of the invention was isolated, and it is about ten fold higher than that of S4 herbicide-resistant mutant tobacco (parent of Hra).

45

10

15

20

25

30

35

*50* 

*55* 

60

## Table 4

Transfer of DNA from Phage Clone 1 to Sensitive N. tabacum Cells

Number of colony forming units derived from 105 protoplast equivalents one month after co-cultivation

	N.t.1	N.t./pl45 <sup>2</sup>
no selection	3.5 x 10 <sup>4</sup>	3.6 x 10 <sup>4</sup>
Kanamycin 50 µg/mL	0	5.9 x 10 <sup>2</sup>
Chlorsulfuron 2 ng/mL	0	0

- 1. Non-cocultured (control) plant cells.
- 2. Plant cells co-cultured with <u>A. tumefaciens</u> harboring pAGS145, kanamycin resistance vector containing the tobacco gene for herbicidesensitive ALS from phage clone 1.

### Table 5

Transfer of DNA from Phage Clone 3 to Sensitive N. tabacum Cells

Number of colony forming units derived from 10<sup>5</sup> protoplast equivalents one month after co-cultivation

	N.t.1	N.t./pl12 <sup>2</sup>	N.t./pl52 <sup>3</sup>
no selection 2:	0 x 10 <sup>4</sup>	$2.0 \times 10^4$	1.6 x 104
Kanamycin 50 μg/mL	0	2:5 x 10 <sup>3</sup>	_
Chlorsulfuron 2 ng/mL	0	0	6.5 x 10 <sup>2</sup>

- 1. Non-cocultured (control) plant cells.
- Plant cells co-cultured with <u>A</u>. <u>tumefaciens</u> harboring pAGS112, kanamycin resistance vector.
- 3. Plant cells co-cultured with <u>A. tumefaciens</u> harboring pAGS152, kanamycin resistance vector containing phage clone 3.

### Table 6

Level of Chlorsulfuron Resistance in Cells of N.tabacum cv. W38 Transformed with Mutant ALS Gene

Number of Colonies Actively Growing After One Month on Selective Medial

chlorsulfuron (ppb)	N. tabacum <sup>2</sup>	N. tabacum/3 mutant ALS gene
0	100	100
20	0	100
50	0	100
200	o ·	100 .
500	0	100 <sup>:</sup>
2000	0	. 99
20000	N.D.4	6 .
50000	N.D.4	0

- One hundred colonies plated at each chlorsulfuron level.
- Colonies derived from non-cocultured (control) plant cells.
- Colonies derived from co-cultivation with <u>A. tumefaciens</u> harboring pAGS152 and initially selected for chlorsulfuron resistance at 2ppb.
- 4. Not determined.

## N. tabacum Culture Media

	•	
Stock	[Final]	Amount/Liter
10%	•	100 mL
100%		10 mL
100%		10 mL
100X		10 mL
1000X		1 mL
1000X		i mL
	0.4M	72.08 gm
	0.4M	136.8
	10X 100X 100X 100X	10X 100X 100X 1000X 1000X

K<sub>3</sub>/S (1) - sucrose. phytohormone regime 1 (elevated)
K<sub>3</sub>/G (1) - glucose. phytohormone regime 1 (elevated)
K<sub>3</sub>/G (2) - glucose. phytohormone regime 2 (reduced)
1 - NAA 3.0 mg/liter 2 - NAA 0.1 mg/liter
BAP 1.0 mg/liter BAP 0.1 mg/liter
bring pH to 5.7, filter sterilize, and
store at 5°.

		•	
C-Medium			
<u>Ingredient</u>	Stock	[Final]	Amount/liter
C-Media majors	10%		100 mL
Fe EDTA	100X	•	10 mL
B5 vitamins	100%		10 mL
MS minors I	1000X		1 mL
MS minors II	1000X	•	1 mL
Mannitol	•	0.2M	36.44 gm
Sucrose		0.1M	34.2 gm
Mes buffer		3.OmM	590 mg
NAA	l mg/mL	0.1 mg/liter	100 ul
BAP	1 mg/mL	0.1 mg/liter	100 ul
BAF	_	to 5.7. filter	sterilize.

and store at 5°.

MSP-Medium (for cell proliferation)				
Ingredient	Stock	[Final]	Amount/liter	
MS majors	10%		100 mL	5
Fe EDTA	100X		10 mL	J
B5 vitamins	100X		10 mL	
MS minors I	1000%		1 mL	10
MS minors II	1000X		1 mL	,,,
Sucrose		0.1 M	34.2 gm	
Mes buffer		3.0 mM	590 mg	15
NAA	1 mg/mL	0.1 mg/liter	100 ul	
BAP	l mg/mL	0.1 mg/liter	100 ul	
	bring pH t	o 5.7, filter s	terilize,	20
	and store	at 5°.		
MSR-Medium (fo	r plant reg	eneration)		25
Ingredient	<u>Stock</u>	[Final]	Amount/liter	
MS major	10X		100 mL	
Fe EDTA	100X		10 mL	30
B5 vitamins	100X		10 mL	
MS minors I	1000X		1 mL	
MS minors II	1000X		1 mL	35
Sucrose		0.1 M	34.2 gm	
Mes buffer		3.0 mM	590 mg	
NAA	1 mg/mL	O.l mg/liter	100 ul	40
BAP	1 mg/mL	1.0 mg/liter	1.0 mL	
bring pH to 5.7,> add agar				
Agar (T.C.)		0.8% (W/V)	8.0 gm	45
autoclave	•			
Kanamycin Sulfate	50 mg/mL	50 ug/mL	1.0 mL	50
or Chlorsulfuron	l mg/mL in 5 mM KOH	as desired	-	
				55

Dispense 25 mL/100x25 mm petri dish and, if required, aseptically add selective agents when media has cooled to  $50^{\circ}$ .

60

	OMS-Medium (for	: plant :	maintenance)	•
	Ingredient	Stock	[Final]	Amount/liter
5	MS majors	10X		100 mL
	Fe EDTA	100X		10 mL
	MS minors I	1000X		1 mL
	MS minors II	1000%	•	1 mL
10	B5 vitamins	100%		10 mL
	Sucrose		3.0% W/V	30 gm
	Mes buffer		3.0 mM	590 mg
15	•	pH 5.7	> add agar	
	Agar (T.C.)		0.8% (W/V)	8.0 gm
<b>20</b> .	autoclave.	dispens	e 50 mL/3" x 4"	Magenta Box

25	Minimal A Medium	•		
25	Ingredient	Stock	[Final]	Amount/liter
	K2HPO4			10.5 g
~	кн <sub>2</sub> Ро			4.5 g
30	(NH) <sub>2</sub> SO <sub>4</sub>			1.0 g
	Sodium citrate 2H <sub>2</sub> O			0.5 g
35	· <b>2</b>	autocla	ave in 990	mL
35	MgSO <sub>4</sub> 7H <sub>2</sub> O	1M	1mM	1.0 mL add
	- 4 2			sterile
40 ·	Glucose	20%		10.0 mL add
				sterile

To solidify media: autoclave agar (15 gm/liter) Difco Bacto. in separate 500 mL volume. Then mix salts and agar before dispensing.

0 257 993

Stock	Ingredient	[Final]	Amount/Liter			
. 14 - (10 <b>V</b> )	NW NO	20.6 mH	16.5 g			
MS major salts (10X)	NH <sup>4</sup> NO <sup>3</sup>	18.8 mH				
	KNO <sub>3</sub> HgSO <sub>4</sub> 7H <sub>2</sub> O	1.5 mH	3.7 g			
•	KH <sub>2</sub> PO <sub>4</sub>		1.7 g			
	CaC1 2H20	3.0 mM	4.4 g			
C-Medium major salts	.NH .NO.	5.0 mM	4.0 g			
	KNO 3	15.0 mM	15.2 g			
(IOA)	HgSO <sub>4</sub> 7H <sub>2</sub> O	3.0 mM	7.4 g			
	KH <sub>2</sub> PO <sub>4</sub>	0.5 mH	0.68 g			
	CaCl <sub>2</sub> 2H <sub>2</sub> O	3.0 mH	4.4 g			
K Medium major salts	KNO,		25.0 g			
(10X)	(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>		1.34 g			
	HgS04 7H20		2.5 g			
	KH <sub>2</sub> PO <sub>4</sub>		2.01 g			
	NH NO3		2.5 g			
	, <b>4 3</b>					
CaCl <sub>2</sub> 2H <sub>2</sub> 0 (100X)	CaCl <sub>2</sub> =2H <sub>2</sub> O		92.3 g			
2 2 2						
Fe-EDTA (100X)	Na PEDTA		3.73 g			
	FeSO <sub>A</sub> -7H <sub>2</sub> O		2.78 g			
(dissolve EDTA entirely before adding FeSO <sub>4</sub> ; pH to 3.0)						
	н <sub>3</sub> во <sub>3</sub>		0.620 g			
Ms minor I (1000X)	MnCl <sub>2</sub> •4H <sub>2</sub> 0		1.980 g			
	Znso <sub>4</sub> •7H <sub>2</sub> o		0.920 g			
	4 2					
MS minor II (1000%)	KI		83 mg			
42 minor, it (1000m)	Na2HoO4-2H2O		25 mg			
	CuSO <sub>4</sub> •5H <sub>2</sub> O		2.5 mg			
	CoCl <sub>2</sub> •6H <sub>2</sub> O		2.16 mg			
	2 2					

<u>Stock</u>	Ingredient	[Final] Amount/Liter
B5 vitemins (100%)	nicotinic acid thiamin HCl pyridoxine HCl myo-inositol	0.1 g 1.0 g 0.1 g 10.0 g
NAA	Naphthelene acetic acid 1 mg/mL (dissolve in dilute KOH	1.0 g
ВАР	Benzylamino  purine l mg/mL  (dissolve in dilute HCl)	1.0 g

## Supplemental Apparatus, Chemicals & Media

MES buffer (2 [N-Morpholino] ethanesulfonic Acid)	Sigma No. M-8250	5
Agarose Type VII Low Gelling Temperature (stock maintained molten at 50°)	Sigma No. A-4018	10
Cellulysin <sup>TM</sup>	Calbiochem 219466	15
Macerase <sup>TM</sup>	Calbiochem 441201	
Cefotaxime, sodium salt dilute w/g.d., sterile H <sub>2</sub> O	Calbiochem 219380	20
store @ 5°, dark, < 10 day as	50 mg/mL stock	
<pre>Kanamycin Sulfate    dilute w/g.d., H<sub>2</sub>O, filter ste    store @ -20°, dark as 50 gm/mI</pre>	Sigma No. K-4000 erile L stock	25
Chlorsulfuron	E. I. du Pont de Nemours and Company, Wilmington, Delaware 19898	30
100 mm x 20 mm tissue culture petri dish	Corning 25020	35
Babcock bottle	Kimble	
Centrifuge (Babcock compatible)	Damon/IEC Division HN-SII	40
Magenta Boxes 3" x 4"	Magenta Corp. 4149 W. Montrose Ave Chicago, IL 68641	4.5
T.C. Agar	KC Biological CR-100	
		50
KDI C II		

#### **EXAMPLE II**

Tobacco DNA from the C3 mutant was prepared and a genomic DNA library in bacteriophage vector EMBL3 was constructed as described in EXAMPLE I. Phage carrying ALS genes were identified, as described in EXAMPLE I, by hybridization to a <sup>32</sup>P-labeled 5' tobacco ALS gene fragment probe.

55

60

65

Six independent recombinant phage were isolated in a screen of 600,000 recombinants from the C3 library. Restriction endonuclease analysis of these isolated phage indicated that the DNA inserts of three phage could be aligned with the SURA gene from the Hra library (phages 35, 36 and 38). The remaining three phage (phage 31, 34 and 37) had DNA inserts corresponding to the SURB gene. It was expected that the ALS gene carried on phages 35, 36 and 38 would be the SURA-C3 gene, encoding herbicide resistant ALS and the ALS gene carried on phages 31, 34 and 37 would be the SURB gene, encoding herbicide sensitive ALS.

DNA fragments from phages 31, 35 and 38 were subcloned into the pUC119 plasmid and subsequently into the pAGS135 binary vector essentially as described in EXAMPLE I. An approximately 8.3 kb Spe I restriction endonuclease fragment from phage 31, analogous to that present in pAGS148 (Fig. 2), but carrying the SURB gene encoding herbicide sensitive ALS, was subcloned in both possible orientations in the vector. An

approximately 6.3 kb Spe I-Sal I restriction endonuclease fragment from phage 35 and an approximately 7.8 kb Spe I-Sal I fragment from phage 38 were subcloned yielding pALS35(ATCC #67424) and pALS38, respectively. The fragments included 2.5 kb in the 5' direction (upstream) of the ALS coding region, 2.0 kb of ALS coding sequence, encoding herbicide resistant enzyme and 1.8 and 3.3 kb, respectively in the 3' direction (downstream) from the ALS coding region. The latter two subcloned fragments contain a BamH I restriction endonuclease site. Partial BamH I digestions or pALS35 and pALS38 were employed for insertion of these plasmids into the BamH I site of the binary vector pAGS135. The ALS genes in the binary vector, designated p312, p313, p351 and p381 (Table 7) were moved into A. tumefaciens by tri-parental mating, as described in EXAMPLE I.

Introduction of the ALS genes into herbicide sensitive tobacco by co-cultivation of plant cells with A. tumefaciens carrying the ALS genes in the binary vector was performed as described in EXAMPLE I. The results of these co-cultivation experiments are shown in Table 7. The ALS gene isolated in phage 31, i.e. the SURB gene encoding herbicide sensitive ALS, yielded no herbicide resistant plant cells, as expected. The ALS gene isolated in phages 35 and 38, i.e. the SURA-C3 gene encoding herbicide resistant ALS did yield herbicide resistant plant cells. Herbicide resistant plant cells arose at a lower frequency than kanamycin resistant plant cells and at a lower frequency than was observed when the SURB-Hra gene was used. This may reflect either the lesser resistance to the herbicide of the ALS enzyme encoded by the SURA-C3 gene compared to that encoded by the SURB-Hra gene, or the lower expression of the SURA-C3 gene compared to the SURB-Hra gene, or both.

- 10

#### Table 7

#### ALS Reintroduction Exp. 3

Transfer of DNA from phage clones 31, 35 & 36 to Sensitive <u>N.tabacum</u> Cells

Number of Colony Forming Units derived from 10<sup>5</sup>
Protoplast Equivalents One Month after Co-cultivation

5

10

40

45

50

*55* 

60

65

	N.t. <sup>1</sup>				W.t./ p351 <sup>5</sup>		15
no selection	2.1x10 <sup>4</sup>	1.5x10 <sup>4</sup>	1.1x10 <sup>4</sup>	1.9x10 <sup>4</sup>	1.5x10 <sup>4</sup>	1.4x10 <sup>4</sup>	20
Kanamycin 50 ug/ml	ο.	88	65	48	139	87	25
Chlorsulfuron 2 ng/ml	. 0	83	0	0	· 32	18	25
1. Non co-cultured pl	lant cel	ls.					30
2. Plant cells co-cul S4/Hra subclone.	ltured w	ith <u>A. t</u>	umefacie	<u>ns</u> harbo	ring pAG	S152.	
3. Plant cells co-cul			umefacie	ns harbo	ring pAG	S312, C3	35

- subclone, φ31 (orientation 1).
  4. Plant cells co-cultured with <u>A. tumefaciens</u> harboring pAGS313, C3
- subclone, \$431 (orientation 2).
- Plant cells co-cultured with <u>A. tumefaciens</u> harboring pAGS351, C3 subclone, φ35.
- Plant cells co-cultured with <u>A. tumefaciens</u> harboring pAGS381, C3 subclone, φ38.

EXAMPLE III

Mutations were made in the wild-type SURA gene of tobacco in vitro in order to make it encode a herbicide resistant ALS. Restriction endonuclease fragments containing part of the SURA gene were subcloned into M13 phage vectors or plasmid vectors containing an M13 origin of replication to allow production of single-stranded DNA. The specific DNA fragment subcloned depended upon the region to be mutagenized in vitro and the availability of restriction endonuclease sites.

Oligonucleotides 16-17 bases in length, which hybridized to the single-stranded DNA from the SURA ALS gene with single base mismatches, were synthesized. These mismatches were designed to convert amino acid codons found in the wild-type ALS gene to the codons found in ALS genes which encode ALS enzymes resistant to sulfonylurea herbicides. These oligonucleotides include 5' GTCAATTGGAGGATC 3', to change trp 591 to leu, 5' GTCAAGTGGCACGTAGG 3', to change pro 197 to ser, 5' ATGTACCTGAGGATATT 3' to change by 5' GAGGTTTGTTGATAGAG 3' to change asp 384 to val, 5'AGGTTTGAGGGATAGAGT 3' to change asp 384 to yal, 5'AGGTTTGAGGGATAGAGT 3' to change asp 384 to glu, 5' TACTGATGATTTCAGG 3' to change ala 205 to asp and 5' CAGGTGGCCCTTCCATG 3' to change ala 122 to pro.

The oligonucleotides were hybridized to single-stranded DNA and used as primers for synthesis of a complementary strand in a reaction catalyzed by the Klenow fragment of DNA polymerase I, following the procedures of Carter et al. (Oligonucleotide site-directed mutagenesis in M13, Anglian Blotechnology Limited, England, 1985). The resulting DNA was transformed into competent E. coli mutL cells (Kramer et al., 1984, Cell 38, 879). Individual plaques or colonies were purified, depending on whether M13 phage vectors (M13mp18/19) or M13 replication origin plasmid vectors (pTZ18R/19R) were used. Mini-preps of single-stranded DNA were made and used in DNA sequencing reactions to Identify clones that carried the mutated bases.

These in vitro constructed site-specific mutations can be incorporated singly or in combination into the wild-type SURA gene which includes the 5' and 3' regulatory sequences needed to provide expression of the gene in plant cells (see EXAMPLE II). This is accomplished by substituting restriction endonuclease fragments carrying the mutations into a plasmid carrying the SURA gene from which the analogous fragment has been removed. The choice of the restriction fragment to substitute depends upon the position of the mutation in the gene and the availability of restriction endonuclease sites. The introduction of the mutated SURA genes into plant cells is then accomplished as described in EXAMPLES I and II. Any of the DNA fragments containing mutations which result in production of herbicide resistant ALS, as disclosed in the description of the invention, can be produced essentially by this method. Furthermore, the mutations need not be made exclusively in the SURA gene. Analogous mutations can be made in the SURB gene or any other plant gene encoding ALS for which DNA sequence information is available.

EXAMPLE IV

20

35

DNA was prepared from Beta vulgaris cv. Sennika (sugarbeet) and a genomic DNA library in bacteriophage lambda vector EMBL3 was constructed as described in EXAMPLE I. Recombinant phage (300,000) were screened by hybridization to a 32P-labeled 5' ALS tobacco gene fragment probe as described in EXAMPLE I. The filters were washed at 42°C (0.1XSSC) and 20 individual clones were isolated. On the second round of purification the recombinant phage were hybridized to both the 3' tobacco ALS gene probe and the 5' probe. In addition, the filters which had been hybridized to the 5' probe were washed at 55°C. Only one clone,  $\phi$ 21, hybridized to both 5' and 3' probes and also remained hybridized after the 55°C wash. Minilysate DNA preparations were made from the 20 clones and digested with EcoR I and Nco I. The different isolates had different restriction endonuclease digestion patterns and again only o21 hybridized to both probes and remained hybridized after a 55°C wash. One phage,  $\phi$ 41, also had a hybridizing band remaining after a 55°C wash but it did not hybridize to the 3' probe. Figure 7 shows the restriction endonuclease map of the phage φ21, together with subclones which have been constructed from it. The ALS coding region was localized to a 3.0 kb BamH I - Hind III fragment by hybridization with 5' and 3' probes from the N. tabacum gene. Both DNA strands of this fragment have been sequenced. The BamH I - Hind III fragment was subcloned into pUC119 of Bluescript vectors; then Exonuclease III or Bal 31 deletions were generated. The dideoxy sequencing method was used. A comparison of the deduced amino acid sequence encoded by the sugarbeet gene with that of the tobacco gene(s) indicates no homology in the first 88 amino acids of the predicted protein (see Figure 8). This region may represent the chloroplast transit peptide. Thereafter the homology is approximately 90% with an insertion of 4 amino acids around residue 290 of tobacco ALS. Inspection of the amino acid residues which define the sites for herbicide resistance identified in tobacco and yeast indicate that these residue are conserved in sugarbeet ALS also. These data allow a straightforward approach to the construction of a gene encoding herbicide resistant sugarbeet ALS enzyme, by site-directed mutagenesis, as described in EXAMPLE

Three sites have been mutagenized in this sugarbeet gene. The codon GCA for ala at position 122 (numbering of amino acid residues from Figure 6) was changed to CCA for pro, the codon CCA for pro at position 197 was changed to GCA for ala and the codon TGG for trp at position 591 was changed to TTG for leu. The double mutation yielding pro to ala at 197 and trp to leu at 591, which mimics the tobacco SURB-Hra gene, was also made by combining the two single mutations.

In order to transform plants with these in vitro constructed mutations in the sugarbeet ALS gene, DNA fragments containing the mutations and extending from the BamH I site about 910 bp in the 5' direction (upstream) of the coding region to the Pst I site about 1000 bp in the 3' direction (downstream) of the coding region (see Figure 17) were cloned into plasmid vector pUC119. These were introduced into the binary vector pAGS140 for transformation into plant cells as described in EXAMPLE I. The binary vector pAGS140 is similar to pAGS135 (Figure 2) except that between the BamH I site and the right border of the T-DNA of the Ti-plasmid of Agrobacterium tumefaciens a gene which will confer resistance on plants to the antibiotic hygromycin was inserted.

Introduction of the ALS genes into herbicide sensitive tobacco and sugarbeet, by co-cultivation of the plant cells with A. tumefaciens carrying the ALS genes in the binary vector, was performed as described in EXAMPLE I and EXAMPLE II. Results of a co-cultivation experiment in tobacco are shown in Table 8. Herbicide resistant transformants were obtained with three of the four mutant sugarbeet ALS genes. The frequency of obtaining herbicide resistant transformants was lower than that for kanamycin resistant transformants, and also lower than the frequency of herbicide resistant transformants obtained when the tobacco SURB-Hra gene was used. It is believed that this results from poor expression of the mutant sugarbeet ALS genes in tobacco. This may reflect either insufficient nucleotide regulatory sequences upstream or downstream of the mutant sugarbeet ALS genes in the DNA fragments used or poor utilization of sugarbeet nucleotide regulatory sequences in tobacco or both. It is expected that the mutant sugarbeet ALS gene which did not yield herbicide resistant transformants would have done so if more transformants had been obtained and tested.

	•	Table B				
. 1 4 -	Daniak					

Herbicide Resistance in Plant Cells Transformed with Site Specific Mutant Sugarbeet ALS Genes

5

10

35

50

55

60

Gene Origin	TODBCCO	Sugarbeet	Sugarbeet	Sugarbeet	<u>Sugarbeet</u>	
Mutation	pro(197)-ala/ trp(591)-leu	pro(197)-ala/ trp(591)-leu	trp(591)-1eu	pro(197)-ala	ala(122)-pro	15
Chlor- sulfuron <sup>R</sup>	698	91	2	35	o	20
Kanamycin <sup>R</sup>	1083	1362	1080	1073	415	25
Ch <sub>1</sub> /Kan <sup>R</sup>	.645	.067	1.85x10 <sup>-3</sup>	.033	-	
	1 ·					30

Gene introductions were done by standard co-cultivation method. For each construction an aliquot of the co-cultured plant cells (2x10<sup>5</sup> starting plant cells) was scored for chlorsulfuron and another aliquot for kanamycin resistance. Selection was with chlorsulforon at 2ppb or kanamycin at 50ppm.

#### EXAMPLE V

The SURB-Hra gene described in EXAMPLE I was transformed into tobacco cultivars by Agrobacterium tumefacters Infection of tobacco leaf disks and progeny of the transformants were analyzed to demonstrate expression of resistance at the whole plant level and inheritance of the herbicide resistance trait. Standard aseptic techniques for the manipulation of sterile media and axenic plant/bacterial cultures were followed, including the use of a laminar flow hood for all transfers. Potted tobacco plants for leaf disk infections were grown in a growth chamber maintained for a 12 hr, 24°C day, 12 hr, 20°C night cycle, with approximately 80% RH, under mixed cool white fluorescent and incandescent lights. Tobacco leaf disk infections were carried out essentially by the method of Horsch, R.B., Fry, J. E., Hoffmann, N. L., Eichholtz, D., Rogers, S.G., Fraley, R.T., (1985). Science 227: 1229-1231.

Young leaves, not fully expanded and approximately 4-6 inches in length, were harvested with a scalpel from approximately 4-6 week old tobacco plants (Nicotiana tabacum CV.NK326 or K14). The leaves were surface sterilized for 30 minutes by submerging them in approximately 500 ml of a 10% Chlorox, 0.1% SDS solution and then rinsed 3 times with sterile deionized water. Leaf disks, 6 mm in diameter, were prepared from whole leaves using a sterile paper punch.

Leaf disks were inoculated by submerging them for several minutes in 20 ml of a 1:10 dilution of an overnight Agrobacterium culture carrying the plasmid pAGS152. Agrobacterium cultures were started by inoculating 10 ml of Min A (EXAMPLE I) broth with a single bacterial colony removed form a Min A plus tetracycline (EXAMPLE VI) plate. The culture was grown for approximately 17-20 hours in 18 mm glass culture tubes in a New Brunswick platform shaker maintained at 28°C.

After inoculation, the leaf disks were placed in petri dishes containing CN agar medium (EXAMPLE VI). The dishes were sealed with parafilm and incubated under mixed fluorescent and "Gro and Sho" plant lights (General Electric) for 2-3 days in a culture room maintained at approximately 25°C.

To rid the leaf disks of Agrobacterium and to select for the growth of transformed tobacco cells, the leaf disks were transferred to fresh CN medium containing 500 mg/l cefotaxime and 100 mg/l kanamycin. Cefotaxime was kept as a frozen 100 mg/ml stock solution and added aseptically (filter sterilized through a 0.45 µm filter) to the media after autoclaving. A fresh kanamycin stock (50 mg/ml) was made for each use and was filter sterilized into the autoclaved media.

Leaf disks were incubated under the growth conditions described above for 3 weeks and then transferred to fresh media of the same composition.

Approximately 1-2 weeks late shoots developing on kanamycin-selected explants were excised with a sterile scalpel and planted in A medium containing 100 mg/l kanamycin. Root formation on selective and non-selective media was recorded within 3 weeks. Shoots which rooted in kanamycin were transferred to soil and grown in a growth chamber as described above. After 3 to 5 weeks, but before flowering had occurred, leaf tissue was excised and used for ALS assays as described in Example VI. The results of these assays, which indicate that a herbicide resistant form of ALS is being produced, are shown in Table 9. The plants exhibiting herbicide resistant ALS activity were then moved to a greenhouse where they were grown to maturity. Individual flowers were bagged to permit self-fertilization without cross-pollination. Mature seeds were harvested and progeny tests were conducted to determine inheritance of the herbicide resistance trait. Seeds were surface sterilized as described above, dried and planted on SG medium (1/4 MS salts, 0.75% sucrose, 0.8% Agar) in the presence or absence of herbicide (DPX-F6025, Classic®). Sensitive seeds germinated, but did not develop further. Results of the progeny analysis are shown in Table 9. A segregration ratio of 3 resistant progeny to 1 sensitive indicated the presence of a single-site insertion of the SURB-Hra gene in the trasnformant which was stably inherited. This was seen in 15 of 17 transformants. Higher ratios of resistant to sensitive progeny indicated multiple insertions at unlinked positions in the genome. The 15/1 ratio indicates the presence of 2 unlinked SURB-Hra genes and 255/1 ratio Indicates 4 unlinked SURB-Hra genes in the transformants K14 #40 and K14 #7, respectively.

Table 10

# ALS Activity of Wild-Type and Transformed Tomato

#### Percent Uninhibited ALS Activity1 1000ppb 100ppb *30* 10ppb Oppb 4 15 5 100 Wild-type 12 25 100 42 Transformant #3 35 26 Transformant #4a. 60 42 100 15 Transformant #4b 29 100 40 25 Transformant #4c 100 58 43 10 29 15 100 Transformant #40

1The ALS activities in each line are relative to the activity in the absence of herbicide which is taken as 100 percent. The sulfonylurea compound used was DPX-F6025, the active ingredient in Classic® herbicide.

55

60

45

50

20

25

EXAMPLE VI

To transform herbicide sensitive tomato to resistance the SURB-Hra gene from tobacco, carried on the binary vector pAGS152 in A. tumefaciens strain LBA4404, was used (see EXAMPLE I). Standard aseptic techniques for the manipulation of sterile media and axenic plant/bacterial cultures were

followed, including the use of a laminar flow hood for all transfers.

Seeds of tomato (Lycopersicon esculentum var. Herbst Red Cherry) was surface sterilized for 30 minutes in a 10% Chlorox, 0.1% SDS solution and rinsed 3 times with sterile deionized water. The seeds were planted in Magenta boxes (Magenta Corp.) containing 100 ml of OMS agar medium and germinated under mixed fluorescent and "Gro and Sho" plant lights (General Electric) in a culture room maintained at approximately 25°C. Cotyledons from 10-15 day old seedlings were used for Agrobacterium inoculation.

Cotyledons were wounded by removing approximately 2 mm of tissue from each end of the cotyledon with a sterile scalpel. Wounded cotyledons were planted in petri dishes on CTM agar medium either with or without 75µM acetosyringone (Aldrich Chemical).

In preparation for the cotyledon inoculation, a single bacterial colony from a Min A + tetracycline (1 µg/ml) agar plate was inoculated into a flask containing 30 ml of Min A broth (EXAMPLE I) and grown for 2 days at 28°C in a New Brunswick platform shaker. On the morning of the cotyledon inoculation, the bacterial culture was diluted with sterile Min A broth to an OD of 0.1 at 650 nM and allowed to multiply to an OD of 0.2 under the growth conditions previously described. This culture was then used undiluted for the inoculation.

CTM agar plates containing the colyledon explants were flooded with 5 ml of the bacterial solution for approximately 5 minutes, before removal of the solution. The plates were then secured with Time Tape (Shamrock Scientific Specialty) on two sides of the dish and incubated for 2 days under mixed fluorescent and "Gro and Sho" plant lights (General Electric) at approximately 25°C for two days.

To rid the plant cultures of Agrobacterium and to select for the growth of transformed tomato cells, the cotyledon explants were transferred to fresh CTM medium containing 500 mg/L cefotaxime and 50 mg/L kanamycin and incubated under the same culture conditions described above for approximately 3 weeks. The cotyledons were than transferred to fresh media of the same composition and selective agents as CTM but with 1/10 the zeatin concentration.

After approximately 2-4 weeks, shoots developing off of kanamycin-selected cotyledons were excised and planted in OMS media containing 500 mg/L cefotaxime and 100 mg/L kanamycin. Tomato shoots which rooted in kanamycin after about 2-3 weeks were transferred to soil in 8° pots and covered with plastic bags. The plants were grown under mixed fluorescent and incandescent lights for a 12 hr, 24°C day; 12 hr, 20°C night cycle, with approximately 80% relative humidity, for one week before removing the plastic bags. The plants were grown for another 2-4 weeks before performing ALS assays. An increase of uninhibited ALS activity in the presence of the sulfonylurea Classic® in leaf extracts from transformed plants was demonstrated in these experiments (Table 10).

# Table 10 ALS Activity of Wild-Type and Transformed Tomato

·		Percei	nt Uninhi	ibited ALS	Activity <sup>1</sup>
		Oppb	10ppb	100ppb	1000ppb
10	Wild-type	100	15	5	4
	Transformant #3	100	42	25	12
15	Transformant #4a.	100	60	42	26
	Transformant #4b	100	29	15	<b>5</b> .
20	Transformant #4c	100	58	43	25
	Transformant #4d	100	29	15	10

The ALS activities in each line are relative to the activity in the absence of herbicide which is taken as 100 percent. The sulfonylurea compound used was DPX-F6025, the active ingredient in Classic® herbicide.

The assay for ALS activity in the absence or presence of herbicide from transformed or untransformed plants was conducted as follows:

1. Grind 2.5 grams of relatively young leaf tissue (4-6 inches in length) in a tissue homogenizer containing 10 ml extraction buffer (100 mM KHPO4) pH 7.5, 0.5 mM MgCl<sub>2</sub>, 10% glycerol, 1 mM pyruvate, to 0.5 mM TPP, 10 nM FAD) and 200 mg Polyclar AT (BDH Blochemicals). Keep on ice.

2. Homogenize extract for approximately 10 seconds in a polytron (Brinkman Instruments) on setting

**#7.** 

3. Centrifuge extract in a Sorvall SS-34 rotor, 20 min. 16K rpm, 4°C.

4. Equilibrate PD-10 (Pharmacia) columns by washing with column buffer (100 mM KHPO<sub>4</sub> pH 7.5, 0.5 mM MgCl<sub>2</sub>, 10% glycerol, 1 mM pyruvate) 5 times.

5. To plant extract supernatant, add cold saturated (NH4)<sub>2</sub>SO<sub>4</sub> to achieve a 50% cut. Incubate on Ice for 30 minutes.

6. Centrifuge extract in SS-34 rotor, 20 minutes, 16K rpm, 4°C. Decant supernatant.

7. Resuspend pellet in 1 ml cold column buffer.

8. Load extract onto column and chase with a volume of column buffer to achieve total volume loaded equal to 2.5 ml. Let this run through column.

9. Elute proteins with 2X volume of extract loaded. Recover in 15 ml Falcon tube placed beneath

10. Set up reaction for each extract in microfuge tubes as follows: 350 μl reaction mix (200 mM pyruvate, 5 mM TPP, 0.9 mM FAD, 5 mM KHPO<sub>4</sub> pH 7.0), 50 μl of either 5 mM KHPO<sub>4</sub> or desired sulfonylurea concentration, and 100 μl plant extract.

11. Incubate reaction for 1 hour, 30°C.

12. To stop reaction, add 50 μl 6 M H<sub>2</sub>SO<sub>4</sub> and incubate at 60°C for 10 minutes. Spin in microfuge 5 minutes.

13. Set up color development tubes as follows: 500  $\mu$ l 0.5% creatin, reaction tube supernatant, 0.5 ml  $\alpha$ -napthol solution (1.5 g  $\alpha$ -napthol in 30 ml 2.5 N NaOH). Mix and heat at 60° C for 20 minutes.

14. Vortex each tube and load 100 µl of each sample into wells of microtiter plate. Read at OD 530.

65

5

25

*30* 

40

45

50

55

## B Callus Induction Medium

<pre>1 package of MS salts (Gibco Murashige    Organics Medium) with 3% sucrose 1 ml of 1 mg/ml NAA 0.2 ml of 1 mg/ml BAP</pre>	per liter pH 5.8	5
0.8% agar		10
<u>CN</u> Shoot Induction Medium	•	
l package of MS salts with 3% sucrose  1 ml of 1 mg/ml NAA  1 ml of 1 mg/ml BAP  0.8% agar	per liter pH 5.8	15
Root Induction Medium		20
<pre>1 package of MS salts (without sucrose) 10 grams sucrose 0.8% agar</pre>	per liter pH 5.8	25
		30
Agrobacterium R-Medium  Add 7.5 g agar to 440 ml H <sub>2</sub> O, autoclave, and keep at 55°C. Add sterile stock 0.5 ml 1 M MgSO <sub>4</sub> 0.5 ml 1 M CaCl <sub>2</sub> 10.0 ml 20% sucrose 5.0 ml 100 mg/ml kanamycin 50.0 ml 10x salts (Na <sub>2</sub> HPO <sub>4</sub> • 7H <sub>2</sub> O 60 g/l; KH <sub>2</sub> PO <sub>4</sub> , 30 g/l; NaCl, 5 g/l;	us of:	35
NH <sub>4</sub> Cl, 10 g/l)  CTM Medium		40
1 pkg MS salts 1 ml B5 vitamins (per 100 ml; Nicotinic Acid 100 mg, thiamine, hydrochlorid hydrochloride 100 mg, M-inositol 10,000 mg) 3 mM MES 3% glucose	de 1000 mg, pyridoxine	45
0.7% agar pH 5.7 Autoclave and add 1 ml 1 mg/ml zeatin stock	•	50
OMS Medium 1 pkg MS salts 1 ml B5 vitamins (see above) 3 mM MES 3% sucrose		55
0.8% agar pH 5.7		
Min A + Tetracycline (1 μg/ml) Medium  1. Add 7.5 g agar to 400 ml H <sub>2</sub> O  2. Make stock:		60

K <sub>2</sub> HPO <sub>4</sub> KH <sub>2</sub> PO <sub>4</sub> (NH <sub>4</sub> ) <sub>2</sub> Sodium	SO <sub>4</sub> Citrate	2H <sub>2</sub> O	5.25 2.25 0.5 0.25	g
			100	ml

10

5

3. Make MgSO<sub>4</sub> • 7H<sub>2</sub>O stock = 20 g/100 ml, autoclaved

4. Make glucose stock - 20% solution, autoclaved

5. Make tetracycline stock - 1.0 mg/ml in ethanol/H<sub>2</sub>O, 50% v/v filter sterilized

To make Min A medium + 1 μg/ml tetracyline:

Mix (1) and (2) 15

Add 0.5 ml of (3), 5 ml of (4), and 0.5 ml of (5)

#### YEB Medium

20		per liter
	Bacto Beef Extract Bacto Yeast Extract	5.0 g 1.0 g
25	Peptone	5.0 g
	Sucrose	5.0 g
	MgSO <sub>4</sub> · 7H <sub>2</sub> O Agar (optional)	0.5 g 15.0 g

*30* 

35

50

Herbicide solutions: A 1 ppm stock solution of sulfonylurea herbicide can be made by dissolving 1 mg of herbicide in 100 ml of 0.01N NH<sub>4</sub>OH, and then diluting 1:10 with 5 mM KHPO<sub>4</sub> pH 7.0. This stock will suffice to assay herbicide concentrations of 100 ppb or lower. If higher concentrations are desired, dissolve 1 mg in 10 mi of 0.01N NH4OH, etc.

Herbicide dilutions: In the standard assay, 50 µl of herbicide is added to 450 µl assay mix and extract, for a 1:10 dilution of herbicide. So, for each concentration to be tested, a 10X solution in 5 mM KHPO<sub>4</sub> pH 7.0 should

be diluted from the stock solution.

**EXAMPLE VII** 

The tobacco SURB-Hra gene encoding herbicide resistant ALS was used to transform Beta vulgaris (sugarbeet) to herbicide resistance by the following Agrobacterium tumefaciens mediated transformation

In order to surface sterilize seeds, 50-100 seeds were placed in a 25 x 100mm sterile petri dish in laminar flow hood and 25-30 ml of 70% ethanol was added. Seeds were agitated by hand 1-2 min., the ethanol was decanted, and 25-30ml 20% Clorox (20 ml commercial bleach/80ml sterile H<sub>2</sub>O/1 drop Tween 80) was added. The seeds were agitated on gyrotary shaker @ 40 rpm for 20 mins., and the bleach was decanted. The bleach sterilization was repeated 2 times for a total of 60 min., and the sterilized seeds were rinsed 3 times for 5 min. each with 25-30ml sterile H2O.

To germinate the seeds, they were plated on 1-2PGo agar solidified medium 12 seed/50 ml media/15x 150mm petri dish, and cultured @ 24°C in darkness.

NOTE: 10-20% contamination of seed may be anticipated for a good, clean seed lot. For this reason seed is plated far apart on large plates and the germinations are monitored continuously and non-contaminated seed are transferred to fresh plates. If contamination is fungal, then transfers are conducted in a laminar flow cabinet with the fan off.

60-80% germination is expected for a good seed lot. Germination is not syncronous. An approximately 14 day period is required to obtain (a) all germinations and (b) sufficient elongation to provide many appropriate

Prepare Agrobacterium overnight (O/N) suspension cultures as described in EXAMPLE I. Freshly plated Agrobacterium have shorter lag times than cultures stored for long periods at 5°C. It is important that log phase bacteria is used as inocula. Therefore, some trial should be conducted to assure an overnight culture which reaches log phase (OD 550 mm 0.4-0.8) before hypocotyl inoculation).

To prepare plant explants, hypocotyls were cut into approximately 0.5cm sections with a sharp surgical blade, and plated immediately onto agar solidified PGo 0.10.1. Do not allow dessication or damage the wound by use of dull blade or by compressing with forceps.

65

To inoculate explants, they were dipped individually into a log phase suspension of the appropriate Agrobacterium strain. They were immersed briefly (1-3 sec.) and arranged in a grid pattern on a fresh plate: 25 explants/100mm plate of agar solidified PGo 0.1/0.1.

The explants were dried by leaving plates open in a laminar flow hood 10-30 min. This concentrates Agrobacteria onto the wound. It also many limit possible damage by water logging. It is important, though, not to dessicate tissue. Close observation of this step is required. The plates were then sealed with parafilm and cultured at 24°C for 3 days.

The explants were collected into liquid PG 0.1/0.1 containing cefotaxime 500µg/ml in a 25 x 100mm petri dish, and shaken gently on a gyrotary shaker at 40 rpm for 1 hr. The media was decanted, replaced with fresh counterselective media, and shaken gently for an additional 2 hrs. The explants were plated in grids, 25/100mm plate agar solidified PG 0.1/0.1 containing cefotaxime 500µg/ml and cultured at 24°C for 3 days.

Selection for transformed plant cells was applied as follows. Explants were transferred to fresh plates of agar solidified PG0.1/0.1 containing vancomycin 100µg/ml or chlorsulfuron, 2ng/ml as selective agents. The number of resistant colonies was counted after 20-30 days. More than one may be observed at each wound. Transformants were excised as follows. Callus was removed from the wound/explant with a surgical blade and cultured independently on fresh selective media. Some Agrobacterium can escape from counter-selection. Additional washes in cefotaxime containing liquid media are possible, as is repeated transfer to cefotaxime containing agar solidified plates. Under the suggested protocol we observed approximately 15% explant contamination, which was an acceptable loss. The results of a transformation experiment using the suggested transformed with the SURB-Hra mutant ALS gene of tobacco, is compared to that of untransformed calli. These results demonstrate that the tobacco SURB-Hra gene, encoding herbicide-resistant ALS, can be expressed efficiently in sugarbeet.

#### Table 11

## Chlorsulfuron (ppb)

30 <u>0</u> 10 <u>30</u> 100 1000 3000 10000 87193/152<sup>1</sup> 15/15 15/15 15/15 15/15 15/15 15/15 3/15 0/15 35 87193/02 15/15 0/15 0/15 nd nđ , nd nd

1Beta vulgaris line 87193 transformed with Agrobacterium tumefaciens LBA4404 carrying plasmid pAGS152.

<sup>2</sup>Untransformed <u>Beta vulgaris</u> line 87193.

55

5

10

20

25

40

45

50

60

## Media and Ammendments

1/2 PG <sub>O</sub>	Ingredient	Stock	(Final)	Amt./Liter
	PG <sub>O</sub> Majors A	10%		50ml
	PGO Majors B	100X		5
	FeEDTA	100X		5
	B5 vitamins	100X		5 .
	MS micronutrients	1000X		ī
	Sucrose		1.5%w/v	15gm
	Mes buffer		3 mM	590mg
	T.C. agar		0.8%W/V	8gn

pH5.7, autoclave sterile, dispense aspectically into 15x150mm petri plates.

PG <sub>0</sub> 0.1/0.1	Ingredient	Stock	(Final)	Amt./Liter
	PG <sub>O</sub> Majors A	10%		100ml
	PGO Majors B	100X	•	10
	FeEDTA	100X		10
	<b>B5 vitamins</b>	100X		10
	MS micronutrients	1000X		1
	Sucrose		3.0%w/v	30gm
	Mes buffer		3 mM	590mg
•	2,4 -D	lmg/ml		100µ1
	Blap	lmg/ml	•	100µ1

pH5.7, autoclave sterile.

PGO 0.1/0.1 agar solidified, as above except with T.C. agar 0.8%w/v. Dispense 25m1/25x100 petri dish.

## Stock Solutions

Stock	Ingredient	ww	(Final)	Amount/: Liter	5
PG <sub>O</sub> Majors A	(10X) KNO <sub>3</sub> (NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub> KC1 MgSO <sub>4</sub> .7H <sub>2</sub> O CaCl <sub>2</sub> .2H <sub>2</sub> O	101.1 132.14 74.55 146.5 147.0	19.8mM 3 8 2	20 4 6 5 3	10
PG <sub>O</sub> Majors B	(100X) NaH <sub>2</sub> PO <sub>4</sub>	120.0	2.lmM ·	25	15
Fe EDTA (100%	C)M FeSO <sub>4</sub> 7H <sub>2</sub> O Na <sub>2</sub> EDTA		100µm 100µm	2.78gm . 3.72	20
	Dissolve EDTA fir dark	st. pH3	.O. Stor	e @ 4°C,	<i>25</i>
B5 vitamins (	100X) Nicotinic acid Thiamine HCl Pyridoxine HCl Myo-inositol		lmg/lit 10 1	0.1gm 1.0 0.1	<i>30</i>
MS micronutri	ents (1000X) MnCl <sub>2</sub> 4H <sub>2</sub> O H <sub>3</sub> BO <sub>3</sub> ZnSO <sub>4</sub> 7H <sub>2</sub> O KI	197.9 01.8 287.5 166	100µm 100 30 5	19800mg 6200 8625 830	35
	NaMoO <sub>4</sub> 2H <sub>2</sub> O CuSO <sub>4</sub> 5H <sub>2</sub> O CoCl <sub>2</sub> 6H <sub>2</sub> O	206 249.7 237.9	1.2 0.1 0.1	250 25 25	40
	Dissolve MnCl <sub>2</sub> 4H Dissolve one at a before adding nex Boil, cool, pH 4.	time an t.	d complet	_	45

## EXAMPLE VIII

The tobacco SURB-Hra gene encoding herbloide resistant ALS was used to transform Brassica napus cv. Olga by the following Agrobacterium tumefaciens mediated transformation procedure.

50

60

65

To surface sterilize seed, they were immersed for 1 min. in 70% ethanol, then 30-60 min. in Clorox/Tween (see EXAMPLE VII). The surface sterilized seeds were germinated on 1/2 MS, 1/2 PGO (SEE EXAMPLE VII), 24°C in the dark, At 5-10 days post germination, hypocotyls were divided into 0.5cm sections and placed on solid I medium containing Acetosyrigone 100 μm (IAS100).

Immediately the explants were dipped individually into a log phase suspension of LBA 4404 containing binary plasmid pAGS15.

The explants were plated back onto IAS100. The <u>Agrobacterium</u> droplet was carefully dried down onto the tissue by leaving the plate open in a laminar flow hood. Co-cultivation was conducted at 24°C in low light or darkness for 3 days.

After 3 days the explants were collected into liquid I medium containing cefotaxime 500mg/L in 100x25mm petri dishes, and shaken on a gyrotory shaker at 40rpm for 3 hrs.

The explants were plated on solid I medium containing cefotaxime 500mg/mL, and cultured for 3 days at 24°C in low light.

The explants were plated on solid I medium containing vancomycln 100 mg/L and kanamycin 100 mg/L. After about 1 month transformed callus appeared as discreet nodules at the ends of explants.

As nodules appeared, they were excised with a sharp scalpel and placed on solid I medium containing kanamycin 100 mg/L.

When transformed callus reached a sufficient size (0.5cm diameter) It was transferred to KR medium containing kanamycin 100 mg/L. This material regenerates fastest if it is plated on fresh media every two weeks. Roots were regenerated on 1/2 MS containing IBA 2µm.

In one experiment, of 100 wound sites (either end of 0.5cm hypocotyl sector) 20 developed callus tissue which was resistant to kanamycin (100 mg/L). Five of the 20 transformed cell lines were subsequently induced to regenerate on kanamycin and somatic siblings for each regenerant genotype were produced by nodal multiplication. These plants were sprayed with various chlorsulfuron concentrations and the results are summarized in Table 12. Two of the five transformants are resistant to chlorsulfuron at levels which are about 10 times greater than that which is lethal to control (untransformed) plants.

Table 12

	0.3 ppm	1 ppm	3 ppm	10 ppm
untransformed	++	+	-	<u>.</u>
R <sub>O</sub> #1	++	++	++	+
R <sub>O</sub> #2	<b>++</b>	+	. <b>-</b>	-
R <sub>O</sub> #3	. ++	+	<b>+</b>	<b>-</b> .
R <sub>O</sub> #4	++	<b>.</b> •	-	-
R <sub>O</sub> #5	+++	++	++	+

<sup>+++</sup> normal growth, no axial induction

₫.

To further demonstrate the expression of the SURB-Hra gene in transformed Brassica napus, an ALS assay in the present of the herbicide chlorsulfuron was performed as described in EXAMPLE VI. The ALS activities of the untransformed parent and transformant R<sub>o</sub> #5 (Table 12) were compared (Table 13). A consistent increase in the percent uninhibited ALS activity was observed in the transformant. Thus, the tobacco SURB-Hra gene, encoding herbicide-resistant ALS, can be expressed in Brassica napus, but that expression is not efficient. Addition of Brassica napus nucleotide regulatory sequences would be expected to increase the level of expression of herbicide resistant ALS and the level of tolerance to foliar application of the herbicide.

60

45

50

5

<sup>++</sup> reduced growth, sublethal at apis, axial induction

<sup>+</sup> reduced growth, lethal at apis, axial induction

<sup>-</sup> lethal

## Table 13

## ALS Activity of Wild Type and Transformed Brassica napus

# Percent Uninhibited ALS Activity1

	Oppb	1ppb	<u>10ppb</u>	100ppb	1000ppb
Wild Type	100.0	86 <sup>.</sup> .6	28.2	10.1	7.6
Transformant Ro#5	100.0	88.1	36.6	20.1	14.5

<sup>1</sup> The ALS activities are relative to that in the absence of herbicide which is taken as 100 percent. The sulfonylurea compound used was chlorsulfuron (DPX-W4189), the active ingredient in Glean® herbicide.

## Brassica napus

## Culture Media

## I Media

Ingredient	Stock	(Final)	Amount/Liter
MS Major Salts MS Micronutrients	10X 1000X		100 ml 1 ml
Fe EDTA	100%	<b>-</b>	1.0 ml
I Vitamins	100%		10 ml
2,4-D	lmg/ml		0.2 ml
Kinetin	lmg/ml	•	3 ml
Sucrose		3% W/V	30 gm
Mannitol		1.8%W/V .	18.2 gm
T.C. agar	•	0.8%w/v	· 8 gm
Mes Buffer		3 mM	0.59 gm

## pH 5.7, autoclave sterile

## KR Media

Ingredient	<b>Stock</b>	(Final)	Amount	/Liter
K3 Major Salts	10X		100	ml
CaCl <sub>2</sub> ZH <sub>2</sub> O	100X		10	ml
MS Micronutrients	1000X		1	ml
Fe EDTA	100X		10	ml
B5 Vitamins	100X		10	ml
Zeatin*	lmg/ml		2	ml
IAA*	lmg/ml		0.1	ml
Sucrose		1% W/V	10	gm
Xylose		0.025%W/V	0.25	gm
Agarose (Type 1, 1d	w E/Eo)	0.25% W/V	. 2.5	gm
Mes Buffer	•	3 mM	0.59	gm

pH 5.7. autoclave sterile

\*\*add these filter sterilized components aseptically

#### Brassica napus

#### Stock Solutions

				•	5
Stock	Ingredient	(Stock)	(Final)	Amount/Liter	
MS Major Salts	nh4no3	10X	20.5 mH	16.5 gm	
	KINO3		18.8	19.0	
	M8504 7H20		1.5	3.7	10
	KH2POA		1.25	1.7	
	CaC12 2H20		3.0	4.4	
K3 Major Salts	KNO3	10X	25.0 mM	25.0 gm	15
	(NH4)2SO4		1.0	1.34	
	MgSOA 7 H20		1.0	2.5	•
	KH2PO4		1.5	2.01	
	ин4и03		3.1	2.5	. 20
CaC1 <sub>2</sub> 2H <sub>2</sub> O	CaCl <sub>2</sub> 2H <sub>2</sub> O	100X	6.3 mH	92.3 gm	
MS Micronutrients	MNCl <sub>2</sub> 4H <sub>2</sub> O	1000X	100 µm	19800 mg	
	H <sub>3</sub> BO <sub>3</sub>		100	6200	25
	ZnSO4 7H20		30	8625	•
	KI		5	830	
	NaHoO4 2H2O		1.2	250	
	CuSO4 5H20		0.1	25	30
	CoCl <sub>2</sub> 6 H <sub>2</sub> O		0.1	25	
Fe EDTA	Na <sub>2</sub> EDTA	100X	100 µm	3.73 gm	
	FeSO <sub>4</sub> 7H <sub>2</sub> O		100	2.78	
					35
I Vitamins	Myo-Inositol	100X	100mg/1	1000mg	
	Thiamine		0.5	50	
	Glycine		2.0	200	
•	Nicotinic acid		5.0	500	40
•	Pyrodoxine		0.5	50	
	Folic acid		0.5	50	
	Biotin		0.05	5	

EXAMPLE IX

The tobacco <u>SURB-Hra</u> gene encoding herbicide resistant ALs was used to transform <u>Cucumis melo</u> cv. Amarillo Oro (melon) to herbicide resistance by the following <u>Agrobacterium tumefaciens mediated</u> transformation procedure. A reference to this procedure is Moreno, V., et al. Plant regeneration from calli of melon. Plant Cell Tissue Organ Culture, 5 (1985) 139-146.

45

50

55

ഞ

65

Surface sterilization of seeds was greatly facilitated by first removing the seed coat. The seeds were then sterilized by rinsing in 70% ethanol for 2 min. then washing in Clorox/Tween (see EXAMPLE VII) for 20 mins. The sterile seeds were washed 3 times in the sterile distilled H<sub>2</sub>O and germinated on OMS at 24°C with a 16 hr. day length.

Cotyledons of 7-14 day old melon seedlings were cut into 5 mm slices with a fresh, sharp scalpel. These explants were dipped into a log phase <u>Argobacterium</u> culture prepared as described in EXAMPLE I, transferred to fresh co-cultivation plates and cultures at 24°C with 16 hr. days for 3 days.

The bacteria were killed by washing the explants for 3 hrs. with gentle agitation in washing media and cultured on fresh selection plates.

The explants were subcultured every 3-4 weeks, dissecting the more compact, darker green sectors away from the white fluffler callus.

When "morphogenic" callus (very dark green, compact, perhaps some recognizable leaves) was seen, it was transferred to regeneration media. The tissue can go directly to shoots instead of going through the

morphogenic stage. Shoots were rooted in rooting media. Approximately 70% of the explants developed callus resistant to kanamycin at 100µm/l. Transformed callus was put on media containing increasing concentrations of chlorsulfuron and growth in the presence of herbicide was determined by weighing the callus after 30 days (Table 14). Some transformants grew as well at high concentrations of chlorsulfuron (1000ppb) as in its absence, e.g. Trans 1, Trans 2 and Trans 7. Thus the tobacco SURB-Hra gene can function to transform melon to high level herbicide resistance.

Table 14 .

10	Chlorsulfuron ppb	Nontransformed	Trans 1	Trans 2	Trans 3	Trans 4	Trans 5
	0	5.7	30.4	30.4	30.4	30.4	30.4
	/ <del>-</del>	0	44.5	19.5	10	14 .	175
	50	Ŏ	25.9	0	7.5	16.6	70
15	100	Ö	51.9	14.8	ο.	0	3.7
	500	Ö	46	26	11.7	10	5.7
	1000	0	19.1	8	28.7	5	0
	2000	0	15.2	18	0	Ö	0
	3000 .	0	41.9	3.3	Ö	0	0
20	4000	-	14.4	0	3.6	Ō	0
	5000	0	14.4	J	3.0		-
•						•	
25			Trans 6	Trans 7	Trans 8	Trans 9	Trans 10
			30.4	26.4	28.3	27.2	40
			28.8	46.5	18.1	27.3	40
	•		18.3	25.5	14.9	11.1	39
<i>30</i>			2	16.1	2	10.4	18.6
			4.7	19.3	2.6	9 .	27.6
			0	19.3	0	8.8	23.5
			0	17.1	10	13.7	17.6
			Ö	20.7	3.4	3	7.6
35			Ö	26.5	7.4	2.6	8.9

Measurements indicate fold increase in weight of callus.

Media OMS MS Salts and Fe EDTA B5 Vitamins 1X Sucrose 3% MES 3mM pH 5.7 0.8% T.C. agar Autoclave 20 min. Basic medium MS Salts and Fe EDTA Myo inositol 100mg/l Thiamine 1mg/l Sucrose 3% MES 3mM pH 5.7 T.C. agar 0.8% Autoclave 20 min. Co-cultivation Medium is Basic medium plus:

40

Acetosyringone 100 μm (acetosyringone is kept as a 100mM stock in DMSO)	
Kinetin 6mg/l	
IAA 1.5 mg/l - Westing Medium in Regio Medium without area about	
Washing Medium is Basic Medium without agar plus: Cefotaxime 500 mg/l	_
Kinetin 6 mg/l	5
IAA 1.5 mg/l	
Selection Medium is Basic Medium plus:	
Kinetin 6 mg/l	
IAA 1.5 mg/l	10
Vancomycin 100 mg/l	10
One of the following selective drugs depending upon Agrobacterium construction:	
Kanamycin 100 mg/l	
Hygromycin 50 mg/l -	
Chlorsulfuron 100 mg/l	15
Regeneration Medium is Basic Medium plus:	
BAP 0.1 mg/i	
Vancomycin 100 mg/l	
Selective drugs as above	
Rooting Medium is OMS plus:	20
IBA 2μm	
Vancomycin 100 mg/l	
Selective drugs as above	
EXAMPLE X	25
The tobacco SURB Hra gene encoding herbicide resistant ALS was used to transform Medicago sativa cv	•
Rangelander (alfalfa) to herbicide resistance by the following Agrobacterium tumefaciens mediated	5 _
transformation procedure. A reference to this procedure is Deak, M., Kiss, G., Koncz, C., and Dudits, D	•
Transformation of Medicago by Agrobacterium mediated gene transfer (preprint).  Plants were grown and subcultured in OMS. The materials used were petioles and stem segments (5mm in	
length) from plants about 2 months from the last subculture.	n <i>30</i>
The petioles and stems of sterile plants were cut into 5mm lengths with a fresh, sharp scalpel. These	
explants were dipped into a log phase Agrobacterium culture prepared as described in EXAMPLE 1	
transferred to fresh co-cultivation plates, and cultured at 24°C with 16 hr. days for 3 days.	•
The bacteria were killed by washing the explants for 3 hrs. with gentle agitation in washing media, and	d 35
cultured on fresh selection plates.	, 35
The explants were subcultured every 3-4 weeks. In about 1 month transformed callus growing out of the	•
wounded ends of the explants was seen, dissected away from the explant and plated on fresh selection media	
When callus became more organized, (areas are darker green and more compact) it was transferred to fresh	
maturation media.	40
When developed embryos appeared, they were transferred to gemination media. After germination, the	
small plants were grown on the same medium.	-
Less than 1% of explants developed callus resistant to kanamycin at 100mg/L. Kanamycin resistant sectors	5
were found to be resistant to the herbicide chlorsulfuron at 50 ppb. Three shoots were produced from	1
kanamycin resistant callus. Tissue from these transformants was assessed for herbicide resistant growth ove	r 45
a range of chlorsulfuron concentrations. One transformant was able to grow at 1000ppb chlorsulfuron; the	3
other two were able to grow at 5000ppb. Thus, the tobacco SURB-Hra gene can function to transform alfalfa to	•
high level herbicide resistance.	
Media .	50
OMS	
MS Saits and Fe EDTA 1X	
B5 Vitamins 1X	
Sucrose 3%	
MES 3mM	55
pH 5.7	
T.C. agar 0.8%	
Autoclave 20 min	
Basic Medium MS Saits and Fe EDTA 1X	
MS Saits and Fe EDTA 1X UM Vitamins 1X *	60
Sucrose 3%	
pH 5.7	
T.C. agar 0.8%	
Autoclave 20 min.	<i>6</i> 5

\*100X UM vitamins (amounts given for 100ml of 100x stock) Thiamine HCI 1g Nicotinic acid 0.5a Pyridoxine HCI 1g Myo inositol 10g Glycine 0.2g Co-Cultivation Medium Basic Medium plus Acetosyringone 100µm (Acetosyringone is kept as a stock of 100mM in DMSO) 10 0.5mg/i 2.4-D BAP 0.2mg/l Washing Medium Basic Medium without agar plus 15 500mg/l Cefotaxime 2.4-D 0.5mg/l BAP 0.2mg/l Selection Medium Basic Medium plus 2.4-D 0.5mg/l BAP 0.2mg/l 100mg/l Vancomycin One of the following selective drugs, depending upon Agrobacterium construction: 25 Hygromycin · 50mg/l Chlorsulfuron 100mg/l Maturation Medium 30 Same as selection medium without 2,4-D Germination Medium Basic Medium plus Vancomycin 100mg/I Selective drugs as above 35 Claims 40 Claims for the following contracting states: DE, GB, FR, IT, NL, SE, CH/LI, BE, LU, and GR 1. A nucleic acid fragment comprising a nucleotide sequence encoding plant acetolactate synthase with increased resistance to one or more sulfonylurea herbicides, said nucleotide sequence comprises at least one sub-sequence which encodes one of the substantially conserved amino acid sub-sequences 45 designated A, B, C, D, E, F, and G in Figure 6, the nucleic acid fragment is further characterized in that at least one of the following conditions is met; a) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence A wherein  $\epsilon_1$  is an amino acid other than alanine, and/or  $\epsilon_2$  is an amino acid other than glycine, b) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence B 50 wherein  $\alpha_1$  is an amino acid other than proline, c) the nucleic acid gragment has a sequence which encodes an amino acid sub-sequence C wherein  $\delta_2$  is an amino acid other than alanine, d) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence D wherein  $\lambda_1$  is an amino acid other than lysine, 55 e) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence E wherein y1 is an amino acid other than aspartic acid, f) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence F wherein β3 is an amino acid other than tryptophan, and/or β8 is an amino acid other than valine and/or  $\beta_7$  is an amino acid other than phenylalanine, and 60 g) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence G wherein  $\sigma_1$  is an amino acid other than methlonine. 2. A nucleic acid fragment according to Claim 1, wherein the fragment has a sequence which encodes an amino acid sub-sequence A having the formula 65 PG<sub>E2</sub>A

56

wherein:	
Pis profine,	
3 is glycine,	
z is an amino acid other than glycine, and	_
A is alanine.	5
3. A nucleic acid fragment according to claim 1 or Claim 2, wherein \$2 is an amino acid selected from the	
group consisting of serine, threonine or cysteine.	
4. A nucleic acid fragment according to Claim 3, wherein $\epsilon_2$ is serine.	
5. A nucleic acid fragment according to Claim 1, wherein the fragment has a sequence which encodes	
an amino acid sub-sequence A having the formula	10
PGGE1	
wherein:	
P is proline,	
G is glycine, and	
e <sub>1</sub> is an amino acid other than alanine or glycine	15
6. A nucleic acid fragment according to Claim 1, wherein the fragment has a sequence which encodes	
an amino acid sub-sequence B having the formula	
GQVa <sub>1</sub>	
wherein:	
G is glycine,	20
Q is glutamine,	
V is valine	
α <sub>1</sub> is an amino acid other than proline.	
7. A nucleic acid fragment according to claim 1 or Claim 6 wherein $\alpha_1$ is an amino acid selected from the	
group consisting of alanine, glycine, arginine, lysine, histidine, serine, cysteine, threonine, glutamine and	25
asparagine.	
8. A nucleic acid fragment according to Claim 7, wherein $\alpha_1$ is alanine, arginine, serine or glutamine.	
9. A nucleic acid fragment according to Claim 1, wherein the fragment has a sequence which encodes	
an amino acid sub-sequence C having the formula	
IGδ₁Dδ₂FQE	<i>30</i>
wherein:	
l is isoleucine,	
G is glycine,	
$\delta_1$ is selected from the group consisting of amino acids,	
D is aspartic acid, and	35
$\delta_2$ is an amino acid other than alanine,	
F is phenylalanine,	
Q is glutamine, and	
E is glutamic acid.	
10. A nucleic acid fragment according to claim 1 or Claim 9, wherein δ2 is an amino acid selected from the	40
group consisting of threonine, serine, cysteine, tyrosine, aspartic acid, glutamic acid, tryptophan,	
histidine, phenylalanine, arginine, and lysine.	
11. A nucleic acid fragment according to Claim 10, wherein $\delta_2$ is threonine, cysteine, aspartic acid,	
glutamic acid, tryptophan or arginine.	
12. A nucleic acid fragment according to Claim 1, wherein the fragment has a sequence which encodes	45
an amino acid sub-sequence D having the formula	
Pλ₁D	
wherein:	
P is proline;	
$\lambda_1$ is an amino acid other than lysine; and	50
D is aspartic acid.	
13. A nucleic acid fragment according to claim 1 or Claim 12, wherein $\lambda_1$ is an amino acid selected from	
the group consisting of glycine, alanine, leucine, isoleucine, valine, threonine, serine, cysteine, tyrosine,	
glutamic acid, aspartic acid, proline, asparagine, glutamine, tryptophan and phenylalanine.	
14. A nucleic acid fragment according to Claim 13, wherein $\lambda_1$ is glycine, leucine, threonine, glutamic	55
acid, aspartic acid, asparagine, tryptophan or proline.	
15. A nucleic acid fragment according to Claim 1, wherein the fragment has a sequence which encodes	
an amino acid sub-sequence G having the formula	
MLGo1HG	
wherein	60
M is methionine,	
Lis leucine,	
G is glycine	
$\sigma_1$ is an amino acid other than methionine,	
His histiding and	65

G is glycine. 16. A nucleic acid fragment according to claim 1 or Claim 15, wherein  $\sigma_1$  is an amino acid selected from the group consisting of glutamic acid, aspartic acid, proline, valine, leucine, or isoleucine. 17. A nucleic acid fragment according to Claim 16, wherein  $\sigma_1$  is proline, glutamic acid, or valine. 18. A nucleic acid fragment according to encodes an amino acid subsequence E having the formula 5 RFDy<sub>1</sub>R wherein: R is arginine, F is phenylalanine, D is aspartic acid, and 10 γ<sub>1</sub> is an amino acid other than aspartic acid. 19. A nucleic acid fragment according to claim 1 or Claim 18, wherein y1 is an amino acid selected from the group consisting of glycine, alanine, valine, leucine, Isoleucine, serine, threonine, cysteine, tyrosine, glutamic acid, proline, asparagine, glutamine, lysine, arginine, tryptophan, histidine, and phenylalanine. 20. A nucleic acid fragment according to Claim 19, wherein γ1 is glycine, valine, serine, glutamic acid, 15 proline, asparagine, lysine, cysteine or tryptophan. 21. A nucleic acid fragment according to Claim 1, wherein the fragment has a sequence which encodes an amino acid sub-sequence F having the formula GB1B8B2QB3B4B5B6B7 20 wherein: G is glycine,  $\beta_1,\beta_2,\beta_3,\beta_4,\beta_5,\beta_6$  and  $\beta_7$  are selected from the group consisting of of amino acids, Q is glutamine, and βs is an amino acid other than valine. 22. A nucleic acid fragment according to claim 1 or Claim 21, wherein  $\beta_1$  is methionine,  $\beta_3$  is tryptophan. 25  $\beta_7$  is phenylalanine and  $\beta_8$  is an amino acid other than valine. 23. A nucleic acid fragment according to any one of Claims 1, 21 and 22 wherein β<sub>8</sub> is an amino acid selected from the group consisting of alanine and glycine. 24. A nucleic acid fragment according to Claim 23 wherein  $\beta_8$  is alanine. 25. A nucleic acid fragment according to Claim 1, wherein the fragment has a sequence which includes 30 an amino acid sub-sequence F having the formula Gβ<sub>1</sub>Vβ<sub>2</sub>Qβ<sub>3</sub>β<sub>4</sub>β<sub>5</sub>β<sub>6</sub>β<sub>7</sub> wherein: G is alvaine.  $\beta_1,\beta_2,\beta_4,\beta_5,\beta_6$  and  $\beta_7$  are selected from the group consisting of amino acids, 35 V is valine. Q is glutamine, and B<sub>3</sub> is an amino acid other than tryptophan or proline. 26. A nucleic acid fragment according to Claim 1, wherein the fragment has a sequence which includes an amino acid sub-sequence F having the formula 40 Gβ<sub>1</sub>Vβ<sub>2</sub>Qβ<sub>3</sub>β<sub>4</sub>β<sub>5</sub>β<sub>6</sub>β<sub>7</sub> wherein: G is glycine,  $\beta_1, \beta_2, \beta_3, \beta_4, \beta_6$  and  $\beta_6$  are selected from the group consisting of amino acids. V is valine, 45 Q is glutamine, and B7 is an amino acid other than phenylalanine. 27. A nucleic acid fragment according to claim 1 or Claim 26 wherein  $\beta_7$  is an amino acid selected from the group consisting of leucine, valine and isoleucine. 28. A nucleic acid fragment according to Claim 27, wherein β<sub>7</sub> is leucine. 50 29. A nucleic acid fragment encoding plant acetolactate synthase which is capable of being incorporated into a nucleic acid construct suitable for transforming a plant containing wild-type acetolactate synthase which is sensitive to a sulfonylurea herbicide compound, said nucleic acid fragment having at least one point mutation relative to the wild-type nucleic acid fragment encoding plant acetolactate synthase such that upon transformation with said nucleic acid construct said plant contains said nucleic acid fragment 55 and exhibits increased resistance to the application of said sulfonylurea herbicide compound. 30. An acetolactate synthase protein encoded by a mucleic acid fragment as claimed in any one of claims 1 to 29.

31. A nucleic acid construct comprising a nucleic acid fragment as claimed in any way of claims 1 to 29.

32. A nucleic acid construct as claimed in claim 31 selected from the cloning vectors deposited at the

33. A nucleic acid construct as claimed in claim 31, wherein a nucleic acid fragment encoding plant acetolactate synthase with increased sulfonylurea herbicide resistance as claimed in any one of claims 1 to 29 is linked to a second nucleic acid fragment conferring a second trait such that when said nucleic acid construct is utilized to transform a plant, the expression of herbicide resistance by said plant upon

ATCC, Rockville, Maryland, U.S.A. under accession numbers 40237, 67124, 67126 and 67424.

60

application of sulfonylurea compounds can be utilized to detect the presence of sald second nucleic acid fragment in sald plant.

34. A microorganism containing a nucleic acid fragment or nucleic acid construct as claimed in any one

of claims 1 to 29 and 31 to 33.

35. A plant containing a nucleic acid fragment or nucleic acid construct as claimed in any one of claims 1 to 29 and 31 to 33.

36. A plant as claimed in claim 35 which is a soybean, maize, sugarbeet, sunflower, tobacco, potato, tomato, alfalfa, melon, cotton, lettuce, wheat or rice plant or a <u>Brassica</u> species.

37. A plant tissue culture transformed with a nucleic acid fragment or nucleic acid construct as claimed in any one of claims 1 to 29 and 31 to 33.

10

20

25

30

35

40

45

50

55

60

65

38. Seed obtained by growing a plant as claimed in claim 35 or claim 36.

39. A method for transforming a plant cell with a nucleic acid fragment as claimed in any one of claims 1 to 29 comprising introducing a nucleic acid construct comprising the nucleic acid fragment into the plant cell.

40. A method for selecting a plant cell transformed with a nucleic acid fragment as claimed in any one of claims 1 to 29 comprising

introducing the nucleic acid fragment into a plant cell whose growth is sensitive to inhibition by herbicides to which the ALS encoded by the nucleic acid fragment is resistant to form a transformed plant cell; and selecting a transformed plant cell whose growth is resistant to inhibition by said herbicides at a selective concentration which inhibits the growth of untransformed cells.

41. A method for controlling the growth of undesired vegetation growing at a locus where a plant transformed with a nucleic acid fragment as claimed in any one of claims 1 to 29 has been cultivated comprising applying to the locus an effective amount of herbicide to which said plant is resistant.

42. A method according to Claim 41 wherein the herbicide is a compound selected from the group consisting of

1

wherein R is H or CH<sub>3</sub>;

J is 
$$R_2$$
 $R_2$ 
 $R_3$ 
 $CH_2$ 
 $S_{R_4}$ 
 $R_4$ 
 $R_5$ 
 $R_6$ 
 $R_6$ 

$$R_8$$
 $P^n$ ;  $R_8$ 
 $P^n$ 
 $P^n$ ;  $P^n$ 
 $P^$ 

R<sub>1</sub> is Cl, Br, NO<sub>2</sub>, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>2</sub>-C<sub>4</sub> alkenyl, CF<sub>3</sub>, C<sub>1</sub>-C<sub>4</sub> alkoxy, C<sub>1</sub>-C<sub>4</sub> haloalkoxy, C<sub>3</sub>-C<sub>4</sub> alkenyloxy, C<sub>2</sub>-C<sub>4</sub> haloalkenyloxy, C<sub>3</sub>-C<sub>4</sub> alkynyloxy, CO<sub>2</sub>R<sub>9</sub>, CONR<sub>10</sub>R<sub>11</sub>, S(O)<sub>m</sub>R<sub>12</sub>, OSO<sub>2</sub>R<sub>12</sub>, phenyl SO<sub>2</sub>N(OCH<sub>3</sub>)CH<sub>3</sub>, SO<sub>2</sub>NR<sub>10</sub>R<sub>11</sub>,

•

į

30

35

40

45

50

55

$$N=N$$
 $R_{21}$ 
 $N-N$ 
 $R_{21}$ 
 $N-N$ 
 $R_{21}$ 
 $N-N$ 
 $R_{21}$ 
 $N-N$ 
 $R_{21}$ 

R<sub>2</sub> is H, Cl, Br, F, CH<sub>3</sub>, NO<sub>2</sub>, SCH<sub>3</sub>, OCF<sub>2</sub>H, OCH<sub>2</sub>CF<sub>3</sub> or OCH<sub>3</sub>;
R<sub>3</sub> is Cl, NO<sub>2</sub>, CO<sub>2</sub>CH<sub>3</sub>, CO<sub>2</sub>C<sub>2</sub>H<sub>5</sub>, SO<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, SO<sub>2</sub>CH<sub>3</sub> or SO<sub>2</sub>C<sub>2</sub>H<sub>5</sub>;
R<sub>4</sub> is C<sub>1</sub>-C<sub>3</sub> alkyl, Cl, Br, NO<sub>2</sub>, CO<sub>2</sub>R<sub>9</sub>, CON(CH<sub>3</sub>)<sub>2</sub>, SO<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, SO<sub>2</sub>N(OCH<sub>3</sub>)CH<sub>3</sub> or S(O)<sub>m</sub>R<sub>12</sub>;

R<sub>5</sub> is C<sub>1</sub>-C<sub>3</sub> alkyl, C<sub>4</sub>-C<sub>5</sub> cycloalkylcarbon, F, Cl, Br, NO<sub>2</sub>, CO<sub>2</sub>R<sub>14</sub>, SO<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, SO<sub>2</sub>R<sub>12</sub> or phenyl;
R<sub>6</sub> is H, C<sub>1</sub>-C<sub>3</sub> alkyl, or CH<sub>2</sub>CH = CH<sub>2</sub>;
R<sub>7</sub> is H, CH<sub>3</sub>, OCH<sub>3</sub>, Cl or Br;
R<sub>8</sub> is H, F, Cl, Br, CH<sub>3</sub>, OCH<sub>3</sub>, CF<sub>3</sub>, SCH<sub>3</sub> or OCF<sub>2</sub>H;
R<sub>9</sub> is C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>3</sub>-C<sub>4</sub> alkenyl or CH<sub>2</sub>CH<sub>2</sub>Cl;

R<sub>10</sub> is H or C<sub>1</sub>-C<sub>3</sub> alkyl;

R <sub>11</sub> is H or C <sub>1</sub> -C <sub>2</sub> alkyl;	
R <sub>12</sub> is C <sub>1</sub> -C <sub>3</sub> alkyt;	
R <sub>13</sub> Is H or CH <sub>3</sub> ;	
$R_{14}$ is $C_1$ - $C_3$ alkyl or $CH_2CH = CH_2$ ;	
m is 0, 1 or 2;	5
n is 1 or 2;	
Q is CH <sub>2</sub> , CHCH <sub>3</sub> or NR <sub>1</sub> s;	
R <sub>15</sub> is H or C <sub>1</sub> -C <sub>4</sub> alkyl;	
Pis O or CH2;	
R <sub>16</sub> is H or CH <sub>3</sub> ;	10
R <sub>17</sub> is C(O)NR <sub>18</sub> R <sub>19</sub> ;	
R <sub>18</sub> is Hor CH <sub>3</sub> ;	
R <sub>19</sub> is CH <sub>3</sub> ;	
R <sub>20</sub> is H, Cl, F, Br, CH <sub>3</sub> , CF <sub>3</sub> , OCH <sub>3</sub> or OCF <sub>2</sub> H;	
R <sub>21</sub> is H or CH <sub>3</sub> ;	15
X is CH <sub>3</sub> , OC <sub>2</sub> H <sub>5</sub> or NHCH <sub>3</sub> ;	,,
Y is CH <sub>3</sub> , C <sub>2</sub> H <sub>5</sub> , OCH <sub>3</sub> , OC <sub>2</sub> H <sub>5</sub> , OCF <sub>2</sub> H, OCH <sub>2</sub> CF <sub>3</sub> , Cl, CH <sub>2</sub> OCH <sub>3</sub> or cyclopropyl;	
Z is CH or N;	
and their agriculturally sultable salts; provided that	
a) when Y is CI, then Z is CH and X is OCH <sub>3</sub> ;	20
b) when Y is OCF <sub>2</sub> H, then Z is CH;	20
c) when J is J-1 and R <sub>1</sub> is OSO <sub>2</sub> R <sub>12</sub> or phenyl, the Y is other than OCF <sub>2</sub> H;	
d) when J is J-2, then Y is other than OCF <sub>2</sub> H or OCH <sub>2</sub> CF <sub>3</sub> ; and	
e) when J is J-3 and R <sub>4</sub> is S(O) <sub>m</sub> R <sub>12</sub> , then Y is other than OCH <sub>2</sub> CF <sub>3</sub> .	
43 . A method according to Claim 41 wherein the herbicide is a compound selected from the group	25
consisting of	
·	
х.	
Y <sub>1</sub> []	
	30
1 7 X	
$\bot$ $\swarrow$ $\swarrow$ $\bigwedge$	
Z, N N SO NAr	
± 2,	
	35
wherein	
Aris	
R	40
The second secon	
K.	
	45
ė i	~~
<b>A</b>	
<sup>®</sup> c	
<b>"</b> c	
Rais C1-C4 alkyl, F. Cl, Br. I. NO2, S(O)pRd, COORe or CF3;	
R <sub>a</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl, F, Cl, Br, I, NO <sub>2</sub> , S(O) <sub>p</sub> R <sub>d</sub> , COOR <sub>e</sub> or CF <sub>3</sub> ; R <sub>b</sub> is H. F. Cl, Br, I, C <sub>1</sub> -C <sub>4</sub> alkyl or COOR <sub>e</sub> ;	E C
Rb Is H, F, Cl, Br, I, C1-C4 alkyl or COORe;	50
Rb Is H, F, Cl, Br, I, C1-C4 alkyl or COORe; Re is H, C1-C4 alkyl, F, Cl, Br, I, CH2ORd, phenyl, NO2 or COORe;	50
R <sub>b</sub> Is H, F, Cl, Br, I, C <sub>1</sub> -C <sub>4</sub> alkyl or COOR <sub>e</sub> ; R <sub>c</sub> is H, C <sub>1</sub> -C <sub>4</sub> alkyl, F, Cl, Br, I, CH <sub>2</sub> OR <sub>d</sub> , phenyl, NO <sub>2</sub> or COOR <sub>e</sub> ; R <sub>d</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl;	50
R <sub>b</sub> Is H, F, Cl, Br, I, C <sub>1</sub> -C <sub>4</sub> alkyl or COOR <sub>e</sub> ; R <sub>c</sub> is H, C <sub>1</sub> -C <sub>4</sub> alkyl, F, Cl, Br, I, CH <sub>2</sub> OR <sub>d</sub> , phenyl, NO <sub>2</sub> or COOR <sub>e</sub> ; R <sub>d</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl; R <sub>e</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl, C <sub>1</sub> -C <sub>4</sub> alkenyl, C <sub>1</sub> -C <sub>4</sub> alkynyl, or 2-ethoxyethyl;	50
R <sub>b</sub> is H, F, Cl, Br, I, C <sub>1</sub> -C <sub>4</sub> alkyl or COOR <sub>e</sub> ; R <sub>c</sub> is H, C <sub>1</sub> -C <sub>4</sub> alkyl, F, Cl, Br, I, CH <sub>2</sub> OR <sub>d</sub> , phenyl, NO <sub>2</sub> or COOR <sub>e</sub> ; R <sub>d</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl; R <sub>e</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl, C <sub>1</sub> -C <sub>4</sub> alkenyl, C <sub>1</sub> -C <sub>4</sub> alkynyl, or 2-ethoxyethyl; V is H, C <sub>1</sub> -C <sub>3</sub> alkyl, allyl, propargyl, benzyl or C <sub>1</sub> -C <sub>3</sub> alkylcarbonyl;	•
R <sub>b</sub> Is H, F, Cl, Br, I, C <sub>1</sub> -C <sub>4</sub> alkyl or COOR <sub>e</sub> ; R <sub>c</sub> is H, C <sub>1</sub> -C <sub>4</sub> alkyl, F, Cl, Br, I, CH <sub>2</sub> OR <sub>d</sub> , phenyl, NO <sub>2</sub> or COOR <sub>e</sub> ; R <sub>d</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl; R <sub>e</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl, C <sub>1</sub> -C <sub>4</sub> alkenyl, C <sub>1</sub> -C <sub>4</sub> alkynyl, or 2-ethoxyethyl; V is H, C <sub>1</sub> -C <sub>3</sub> alkyl, allyl, propargyl, benzyl or C <sub>1</sub> -C <sub>3</sub> alkylcarbonyl; X <sub>1</sub> , Y <sub>1</sub> , and Z <sub>1</sub> , are independently H, F, Cl, Br, I, C <sub>1</sub> -C <sub>4</sub> alkyl, C <sub>1</sub> -C <sub>2</sub> alkylthlo or C <sub>1</sub> -C <sub>4</sub> alkoxy; and	50 55
R <sub>b</sub> Is H, F, Cl, Br, I, C <sub>1</sub> -C <sub>4</sub> alkyl or COOR <sub>e</sub> ; R <sub>c</sub> is H, C <sub>1</sub> -C <sub>4</sub> alkyl, F, Cl, Br, I, CH <sub>2</sub> OR <sub>d</sub> , phenyl, NO <sub>2</sub> or COOR <sub>e</sub> ; R <sub>d</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl; R <sub>e</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl, C <sub>1</sub> -C <sub>4</sub> alkenyl, C <sub>1</sub> -C <sub>4</sub> alkynyl, or 2-ethoxyethyl; V is H, C <sub>1</sub> -C <sub>3</sub> alkyl, allyl, propargyl, benzyl or C <sub>1</sub> -C <sub>3</sub> alkylcarbonyl; X <sub>1</sub> , Y <sub>1</sub> , and Z <sub>1</sub> , are independently H, F, Cl, Br, I, C <sub>1</sub> -C <sub>4</sub> alkyl, C <sub>1</sub> -C <sub>2</sub> alkylthlo or C <sub>1</sub> -C <sub>4</sub> alkoxy; and p is 0, 1 or 2.	•
R <sub>b</sub> is H, F, Cl, Br, I, C <sub>1</sub> -C <sub>4</sub> alkyl or COOR <sub>6</sub> ; R <sub>c</sub> is H, C <sub>1</sub> -C <sub>4</sub> alkyl, F, Cl, Br, I, CH <sub>2</sub> OR <sub>d</sub> , phenyl, NO <sub>2</sub> or COOR <sub>6</sub> ; R <sub>d</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl; R <sub>e</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl, C <sub>1</sub> -C <sub>4</sub> alkenyl, C <sub>1</sub> -C <sub>4</sub> alkynyl, or 2-ethoxyethyl; V is H, C <sub>1</sub> -C <sub>3</sub> alkyl, allyl, propargyl, benzyl or C <sub>1</sub> -C <sub>3</sub> alkylcarbonyl; X <sub>1</sub> , Y <sub>1</sub> , and Z <sub>1</sub> , are independently H, F, Cl, Br, I, C <sub>1</sub> -C <sub>4</sub> alkyl, C <sub>1</sub> -C <sub>2</sub> alkylthlo or C <sub>1</sub> -C <sub>4</sub> alkoxy; and p is 0, 1 or 2.  44. A method according to Claim 41 wherein the herbicide is a compound selected from the group	•
R <sub>b</sub> Is H, F, Cl, Br, I, C <sub>1</sub> -C <sub>4</sub> alkyl or COOR <sub>e</sub> ; R <sub>c</sub> is H, C <sub>1</sub> -C <sub>4</sub> alkyl, F, Cl, Br, I, CH <sub>2</sub> OR <sub>d</sub> , phenyl, NO <sub>2</sub> or COOR <sub>e</sub> ; R <sub>d</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl; R <sub>e</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl, C <sub>1</sub> -C <sub>4</sub> alkenyl, C <sub>1</sub> -C <sub>4</sub> alkynyl, or 2-ethoxyethyl; V is H, C <sub>1</sub> -C <sub>3</sub> alkyl, allyl, propargyl, benzyl or C <sub>1</sub> -C <sub>3</sub> alkylcarbonyl; X <sub>1</sub> , Y <sub>1</sub> , and Z <sub>1</sub> , are independently H, F, Cl, Br, I, C <sub>1</sub> -C <sub>4</sub> alkyl, C <sub>1</sub> -C <sub>2</sub> alkylthlo or C <sub>1</sub> -C <sub>4</sub> alkoxy; and p is 0, 1 or 2.	•
R <sub>b</sub> is H, F, Cl, Br, I, C <sub>1</sub> -C <sub>4</sub> alkyl or COOR <sub>6</sub> ; R <sub>c</sub> is H, C <sub>1</sub> -C <sub>4</sub> alkyl, F, Cl, Br, I, CH <sub>2</sub> OR <sub>d</sub> , phenyl, NO <sub>2</sub> or COOR <sub>6</sub> ; R <sub>d</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl; R <sub>e</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl, C <sub>1</sub> -C <sub>4</sub> alkenyl, C <sub>1</sub> -C <sub>4</sub> alkynyl, or 2-ethoxyethyl; V is H, C <sub>1</sub> -C <sub>3</sub> alkyl, allyl, propargyl, benzyl or C <sub>1</sub> -C <sub>3</sub> alkylcarbonyl; X <sub>1</sub> , Y <sub>1</sub> , and Z <sub>1</sub> , are independently H, F, Cl, Br, I, C <sub>1</sub> -C <sub>4</sub> alkyl, C <sub>1</sub> -C <sub>2</sub> alkylthlo or C <sub>1</sub> -C <sub>4</sub> alkoxy; and p is 0, 1 or 2.  44. A method according to Claim 41 wherein the herbicide is a compound selected from the group	•
R <sub>b</sub> is H, F, Cl, Br, I, C <sub>1</sub> -C <sub>4</sub> alkyl or COOR <sub>6</sub> ; R <sub>c</sub> is H, C <sub>1</sub> -C <sub>4</sub> alkyl, F, Cl, Br, I, CH <sub>2</sub> OR <sub>d</sub> , phenyl, NO <sub>2</sub> or COOR <sub>6</sub> ; R <sub>d</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl; R <sub>e</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl, C <sub>1</sub> -C <sub>4</sub> alkenyl, C <sub>1</sub> -C <sub>4</sub> alkynyl, or 2-ethoxyethyl; V is H, C <sub>1</sub> -C <sub>3</sub> alkyl, allyl, propargyl, benzyl or C <sub>1</sub> -C <sub>3</sub> alkylcarbonyl; X <sub>1</sub> , Y <sub>1</sub> , and Z <sub>1</sub> , are independently H, F, Cl, Br, I, C <sub>1</sub> -C <sub>4</sub> alkyl, C <sub>1</sub> -C <sub>2</sub> alkylthlo or C <sub>1</sub> -C <sub>4</sub> alkoxy; and p is 0, 1 or 2.  44. A method according to Claim 41 wherein the herbicide is a compound selected from the group	•

wherein

III

1

ŧ

2

or x<sub>3</sub> COOR<sub>1</sub>

25

40

45

50

55

60

5

10

15

Rt is C1-C4 alkyl;

Rg is C1-C4 alkyl or C3-C6 cycloalkyl;

30 A<sub>1</sub> is COOR<sub>1</sub>, CH<sub>2</sub>OH or CHO;

R<sub>I</sub> is H; C<sub>1</sub>-C<sub>12</sub> alkyl optionally substituted by C<sub>1</sub>-C<sub>3</sub> alkyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl or phenyl; C<sub>3</sub>-C<sub>5</sub> alkenyl optionally substituted by phenyl or 1-2 C<sub>1</sub>-C<sub>3</sub> alkyl, F, Cl, Br or I; or C<sub>3</sub>-C<sub>5</sub> alkynyl optionally substituted by phenyl or 1-2 C<sub>1</sub>-C<sub>3</sub> alkyl, F, Cl, Br or I;

B is H; C(O)C1-C6 alkyl or C(O)phenyl optionally substituted by Cl, NO2 or OCH3;

35 X2 is H, F, Cl, Br, I, OH or CH3;

Y<sub>2</sub> and Z<sub>2</sub> are independently H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, F, Cl, Br, I, phenyl, NO<sub>2</sub>, CN, CF<sub>3</sub> or SO<sub>2</sub>CH<sub>3</sub>;

X<sub>3</sub> is H, C<sub>1</sub>-C<sub>3</sub> alkyl, F, Cl, Br, I or NO<sub>2</sub>; and

L. M. Q and Rh are independently H, F, Cl, Br, I, CH<sub>3</sub>, OCH<sub>3</sub>, NO<sub>2</sub>, CF<sub>3</sub>, CN, N(CH<sub>3</sub>)<sub>2</sub>, NH<sub>2</sub>, SCH<sub>3</sub> or SO<sub>2</sub>CH<sub>3</sub> provided that only one of M or Q may be a substituent other than H, F, Cl, Br, I, CH<sub>3</sub> or OCH<sub>3</sub>.

Claims for the following contracting states: AT and ES

- 1. A method of preparing a nucleic acid fragment comprising a nucleotide sequence encoding plant acetolactase synthase with increased resistance to one or more sulfonylurea herbicides, wherein a nucleotide sequence encoding plant acetolactate synthase and comprising at least one sub-sequence which encodes one of the substantially conserved amino acid sub-sequences designated A,B,C,D,E,F and G in Figure 6 is mutated such that at least one of the following conditions is met:
  - a) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence A wherein  $\epsilon_1$  is an amino acid other than alanine, and/or  $\epsilon_2$  is an amino acid other than glycine,
  - b) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence B wherein  $\alpha_1$  is an amino acid other than proline,
  - c) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence C wherein  $\delta_z$  is an amino acid other than alanine,
  - d) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence D wherein  $\lambda_1$  is an amino acid other than lysine,
  - e) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence E wherein γ<sub>1</sub> is an amino acid other than aspartic acid.
  - f) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence F wherein  $\beta_3$  is an amino acid other than tryptophan, and/or  $\beta_8$  is an amino acid other than valine and/or  $\beta_7$  is an amino acid other than phenylalanine, and
  - g) the nucleic acid fragment has a sequence which encodes an amino acid sub-sequence G wherein on is an amino acid other than methionine.
- A method according to Claim 1, wherein the fragment has a sequence which encodes an amino acid sub-sequence A having the formula

PG<sub>2</sub>A

65 wherein:

P is proline,	
G is glycine,	
ε2 is an amino acid other than glycine, and	
A is alanine.  3. A method according to claim 1 or Claim 2, wherein $\epsilon_2$ is an amino acid selected from the group	5
consisting of sedne, threonine or cysteine.	•
4. A method according to Claim 3, wherein ε₂ is serine.	
5. A method according to Claim 1, wherein the fragment has a sequence which encodes an amino acid	
sub-sequence A having the formula	
PGGE1	10
wherein:	
P is proline,	
G is glycine, and	
ɛ1 is an amino acid other than alanine or glycine.	
6. A method according to Claim 1, wherein the fragment has a sequence which encodes an amino acid	15
sub-sequence B having the formula	
GQVa1	
wherein:	
G is glycine,	
Q is glutamine,	20
V is valine	
$lpha_1$ is an amino acid other than proline.	
7. A method according to claim 1 or Claim 6, wherein $\alpha_1$ is an amino acid selected from the group	
consisting of alanine, glycine, arginine, lysine, histidine, serine, cysteine, theonine, glutamine and	
asparagine.	25
8. A method according to Claim 7, wherein a1 is alanine, arginine, serine or glutamine.	
<ol><li>A method according to Claim 1, wherein the fragment has a sequence which encodes an amino acid</li></ol>	
sub-sequence C having the formula	
IG8₁D8₂FQE	
wherein:	30
I is isoleucine,	
G is glycine,	
81 is selected from the group consisting of amino acids,	
D is aspartic acid, and δ <sub>2</sub> is an amino acid other than alanine.	35
10. A method according to claim 1 or Claim 9 wherein $\delta_2$ is an amino acid selected from the group	33
consisting of threonine, serine, cysteine, tyrosine, aspartic acid, glutamic acid, tryptophan, histidine,	
phenylalanine, arginine, and lysine.	
11. A method according to Claim 10, wherein δ <sub>2</sub> is threonine, cysteine, aspartic acid, glutamic acid,	•
tryptophan or arginine.	40
12. A method according to Claim 1, wherein the fragment has a sequence which encode an amino acid	
sub-sequence D having the formula	
Pλ <sub>1</sub> D	
wherein:	
P is proline:	45
λ <sub>1</sub> is an amino acid other than lysine; and	
D is aspartic acid.	
13. A method according to claim 1 or Claim 12, wherein $\lambda_1$ is an amino acid selected from the group	
consisting of glycine, alanine, leucine, isoleucine, valine, threonine, serine, cysteine, tyrosine, glutamic	
acid, aspartic acid, proline, asparagine, glutamine, tryptophan and phenylalanine.	50
14. A method according to Claim 13, wherein λ <sub>1</sub> is glycine, leucine, threonine, glutamic acid, aspartic	
acid, asparagine, tryptophan or proline.  15. A method according to Claim 1, wherein the fragment has a sequence which encodes an amino acid	
sub-sequence G having the formula	55
MLGo1HG wherein	٠.
Mis methionine,	
Lis leucine.	
G is glycine	
σ <sub>1</sub> is an amino acid other than methionine,	60
H is histidine, and	
G is glycine.	
16. A method according to claim 1 or Claim 15, wherein σ <sub>1</sub> is an amino acid selected from the group	
consisting of glutamic acid, aspartic acid, proline, valine, leucine, or isoleucine.	
17. A method according to Claim 16, wherein a is proline glutamic acid, or valine	64

18. A method according to Claim 1, wherein the fragment has a sequence which encodes an amino acid

sub-sequence E having the formula RFDy<sub>1</sub>R wherein: R is arginine, F is phenylalanine, D is aspartic acid, and  $\gamma_1$  is an amino acid other than aspartic acid. 19. A method according to claim 1 or Claim 18, wherein  $\gamma_1$  is an amino acid selected from the group consisting of glycine, alanine, valine, leucine, isoleucine, serine, threonine, cysteine, tyrosine, glutamic 10 acid, proline, asparagine, glutamine, lysine, arginine, tryptophan, histidine, and phenylalanine. 20. A method according to Claim 19, wherein  $\gamma_1$  is glycine, valine, serine, glutamic acid, proline, asparagine, lysine, cysteine or tryptophan. 21. A method according to Claim 1, wherein the fragment has a sequence which encodes an amino acid sub-sequence F having the formula 15 Gβ<sub>1</sub>β<sub>8</sub>β<sub>2</sub>Qβ<sub>3</sub>β<sub>4</sub>β<sub>5</sub>β<sub>6</sub>β<sub>7</sub> wherein: G is glycine,  $\beta_1,\beta_2,\beta_3,\beta_4,\beta_5,\beta_6$  and  $\beta_7$  are selected from the group consisting of of amino acids, Q is glutamine, and 20 β<sub>8</sub> is an amino acid other than valine. 22. A method according to claim 1 or Claim 21, wherein  $\beta_1$  is methionine,  $\beta_3$  is tryptophan,  $\beta_7$  is phenylalanine and  $\beta_8$  is an amino acid other than valine. 23. A method according to any one of Claims 1, 21 and 22 wherein  $\beta_{\delta}$  is an amino acid selected from the group consisting of alanine and glycine. 25 24. A method according to Claim 23 wherein β<sub>8</sub> is alanine. 25. A method according to Claim 1, wherein the fragment has a sequence which includes an amino acid sub-sequence F having the formula Gβ<sub>1</sub>Vβ<sub>2</sub>Qβ<sub>3</sub>β<sub>4</sub>β<sub>5</sub>β<sub>6</sub>β<sub>7</sub> wherein: 30 G is glycine,  $\beta_1,\beta_2,\beta_4,\beta_5,\beta_6$  and  $\beta_7$  are selected from the group consisting of amino acids, V is valine. Q is glutamine, and  $\beta_3$  is an amino acid other than tryptophan or proline. 35 26. A nucleic acid fragment according to Claim 1, wherein the fragment has a sequence which includes an amino acid sub-sequence F having the formula Gβ<sub>1</sub>Vβ<sub>2</sub>Qβ<sub>3</sub>β<sub>4</sub>β<sub>5</sub>β<sub>6</sub>β<sub>7</sub> wherein: G is glycine, 40  $\beta_1, \beta_2, \beta_3, \beta_4, \beta_5$  and  $\beta_6$  are selected from the group consisting of amino acids, V is valine, Q is glutamine, and By is an amino acid other than phenylalanine. 27. A method according to claim 1 or Claim 26 wherein  $\beta_7$  is an amino acid selected from the group 45 consisting of leucine, valine and isoleucine. 28. A method according to Claim 27 wherein  $\beta_7$  is leucine. 29. A method of preparing a nucleic acld fragment encoding plant acetolactate synthase with increased resistance to one or more sulfonylurea herbicides wherein at least one point mutation is introduced at an appropriate site in a wild-type nucleotide sequence encoding plant acetolactate synthase. 50 30. A method of preparing a nucleic acid construct comprising a nucleotide sequence encoding plant acetolactate synthase with increased resistance to one or more sulfonylurea herebicides which comprises linking a nucleic acid fragment prepared by a method as claimed in any one of claims 1 to 29 to a second nucleic acid fragment. 31. A method as claimed in claim 30 wherein said second nucleic acid fragment confers a second trait 55 such that, when said nucleic acid construct is ultilized to transform a plant, the expression of herbicide resistance by said plant upon application of sulfonylurea compounds can be utilized to detect the presence of said second nucleic acid fragment in said plant. 32. A nucleic acid construct selected from the cloning vectors deposited at the ATCC, Rockville, Maryland, U.S.A under accession numbers 40237, 67124, 67126 and 67424. 60 33. A method of preparing a plant acetolactate synthase with increased resistance to one or more sulfonyturea herbicides, which comprises expressing a plant acetolactate synthase-encoding sequence within a nucleic acid fragment or nucleic acid construct as defined in any one of claims 1 to 32. 34. A microorganism containing a nucleic acid fragment or nucleic acid construct as defined in any one of claims 1 to 32. 65

35. A plant containing a nucleic acid fragment or nucleic acid construct as defined in any one of claims 1 to 32. 36. A plant as claimed in claim 35 which is a soybean, maize sugarbeet, sunflower, tobacco, potato, tomato, affalfa, melon, cotton, lettuce, wheat or rice plant or a Brassica species. 37. A plant tissue culture transformed with a nucleic acid fragment or nucleic acid construct as defined in any one of claims 1 to 32. 38. Seed obtained by growing a plant as claimed in claim 35 or claim 36. 39. A method for transforming a plant cell with a nucleic acid fragment as defined in any one of claims 1 to 29 comprising introducing a nucleic acid construct comprising the nucleic acid fragment into the plant · 10 40. A method for selecting a plant cell transformed with a nucleic acid fragment as defined in any one of claims 1 to 29 comprising introducing the nucleic acid fragment into a plant cell whose growth is sensitive to inhibition by herbicides to which the ALS encoded by the nucleic acid fragment is resistant to form a transformed plant cell; and selecting a transformed plant cell whose growth is resistant to inhibition by said herbicides at a selective 15 concentration which inhibits the growth of untransformed cells. 41. A method for controlling the growth of undesired vegetation growing at a locus where a plant transformed with a nucleic acid fragment as defined in any one of claims 1 to 29 has been cultivated comprising applying to the locus an effective amount of herbicide to which said plant is resistant. 42. A method according to Claim 41 wherein the herbicide is a compound selected from the group consisting of 25 30 I wherein 35 R is H or CH3; 40 45 50 60

J is 
$$R_2$$
 $R_2$ 
 $R_3$ 
 $CH_2$ 
 $SR_4$ 

10

 $R_5$ 
 $R_7$ 
 $R_5$ 
 $R_6$ 
 $R_6$ 
 $R_6$ 
 $R_6$ 
 $R_6$ 
 $R_6$ 
 $R_6$ 
 $R_6$ 
 $R_7$ 
 $R_7$ 
 $R_8$ 
 $R_8$ 

$$R_8$$
 $\stackrel{H}{\longrightarrow}$ 
 $\stackrel{\circ}{\longrightarrow}$ 
 $\stackrel{\circ}{\longrightarrow}$ 

 $R_1$  is Cl, Br, NO2, C1-C4 alkyl, C2-C4 alkenyl, CF3, C1-C4 alkoxy, C1-C4 haloalkoxy, C3-C4 alkenyloxy, C2-C4 haloalkenyloxy, C3-C4 alkynyloxy, C02R9, C0NR10R11, S(0)\_mR12, OSO2R12 , phenyl SO2N(OCH3)CH3, SO2NR10R11,

đ

4

ŝ

Š

30

35

40

55

R<sub>2</sub> is H, Cl, Br, F, CH<sub>3</sub>, NO<sub>2</sub>, SCH<sub>3</sub>, OCF<sub>2</sub>H, OCH<sub>2</sub>CF<sub>3</sub> or OCH<sub>3</sub>;
R<sub>3</sub> is Cl, NO<sub>2</sub>, CO<sub>2</sub>CH<sub>3</sub>, CO<sub>2</sub>C<sub>2</sub>H<sub>5</sub>, SO<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, SO<sub>2</sub>CH<sub>3</sub> or SO<sub>2</sub>C<sub>2</sub>H<sub>5</sub>;
R<sub>4</sub> is C<sub>1</sub>-C<sub>3</sub> alkyl, Cl, Br, NO<sub>2</sub>, CO<sub>2</sub>R<sub>9</sub>, CON(CH<sub>3</sub>)<sub>2</sub>, SO<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, SO<sub>2</sub>N(OCH<sub>3</sub>)CH<sub>3</sub> or S(O)<sub>m</sub>R<sub>12</sub>;
R<sub>5</sub> is C<sub>1</sub>-C<sub>3</sub> alkyl, C<sub>4</sub>-C<sub>5</sub> cycloalkylcarbonyl, F, Cl, Br, NO<sub>2</sub>, CO<sub>2</sub>R<sub>14</sub>, SO<sub>2</sub>N(CH<sub>3</sub>)<sub>2</sub>, SO<sub>2</sub>R<sub>12</sub> or phenyl;
R<sub>6</sub> is H, C<sub>1</sub>-C<sub>3</sub> alkyl, or CH<sub>2</sub>CH = CH<sub>2</sub>;
R<sub>7</sub> is H, CH<sub>3</sub>, OCH<sub>3</sub>, Cl or Br;
R<sub>8</sub> is H, F, Cl, Br, CH<sub>3</sub>, OCH<sub>3</sub>, CF<sub>3</sub>, SCH<sub>3</sub> or OCF<sub>2</sub>H;
R<sub>9</sub> is C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>3</sub>C<sub>4</sub> alkenyl or CH<sub>2</sub>CH<sub>2</sub>Cl;
R<sub>10</sub> is H or C<sub>1</sub>-C<sub>3</sub> alkyl;

R <sub>11</sub> is H or C <sub>1</sub> -C <sub>2</sub> alkyl;	
R <sub>12</sub> is C <sub>1</sub> -C <sub>3</sub> alkyl;	
R <sub>13</sub> is Hor CH <sub>3</sub> ;	
R14 is C1-C3 alkyl or CH2CH = CH2;	5
mis 0, 1 or 2;	
n is 1 or 2; Q is CH <sub>2</sub> , CHCH <sub>3</sub> or NR <sub>15</sub> ;	
115 CH2; CHCH3 CH 4415. R <sub>15</sub> is H or C₁-C₄ alkyl;	
Pis O or CH <sub>2</sub> ;	
R <sub>16</sub> is H or CH <sub>3</sub> ;	10
R <sub>17</sub> is C(O)NR <sub>18</sub> R <sub>19</sub> :	
R <sub>18</sub> is H or CH <sub>3</sub> ;	
R <sub>18</sub> Is CH <sub>3</sub> ;	
R <sub>20</sub> is H, Cl, F, Br, CH <sub>3</sub> , CF <sub>3</sub> , OCH <sub>3</sub> or OCF <sub>2</sub> H;	
R21 Is H or CH3:	15
X is CH3: OCH3, OC2H5 or NHCH3;	
Y Is CH <sub>3</sub> , C <sub>2</sub> H <sub>5</sub> , OCH <sub>3</sub> , OC <sub>2</sub> H <sub>5</sub> , OCF <sub>2</sub> H, OCH <sub>2</sub> CF <sub>3</sub> , CI, CH <sub>2</sub> OCH <sub>3</sub> or cyclopropyl;	
Z is CH or N:	
and their agriculturally suitable saits; provided that	~~
al William Lis Ci, men Lis Chand Xis Cons.	20
b) when Y is OCF <sub>2</sub> H, then Z is CH;	
c) when J is J-1 and R <sub>1</sub> is OSO <sub>2</sub> R <sub>12</sub> or phenyl, then Y is other than OCF <sub>2</sub> H;	
d) when J is J-2, then Y is other than OCF <sub>2</sub> H or OCH <sub>2</sub> CF <sub>3</sub> ; and	
e) when J is J-3 and R <sub>4</sub> is S(O) <sub>m</sub> R <sub>12</sub> , then Y is other than OCH <sub>2</sub> CF <sub>3</sub> .  43. A method according to Claim 41 wherein the herbicide is a compound selected from the group	25
consisting of	
·	
v X <sub>1</sub>	
1  N	<i>30</i>
Y	
Z <sub>1</sub> N N SO <sub>2</sub> NAr	
1 2.	
V	35
wherein	
Aris	
<b>.</b>	40
Kb m	•
<sup>R</sup> ♣	
$\checkmark$	45
R_	
2007 - 2007	
R <sub>B</sub> is C <sub>1</sub> -C <sub>4</sub> alkyl, F, Cl, Br, I, NO <sub>2</sub> , S(O) <sub>p</sub> R <sub>d</sub> , COOR <sub>e</sub> or CF <sub>3</sub> ;	
Rb is H, F, Cl, Br, I, C1-C4 alkyl or COORe;	50
Rc is H, C1-C4 alkyl, F, Cl, Br, I, CH2ORd, phenyl, NO2 or COORe;	
Rd is C1-C4 alkyl;	
Re is C1-C4 alkyl, C1-C4 alkenyl, C1-C4 alkynyl, or 2-ethoxyethyl;	
V is H, C1-C3 alkyl, allyl, propargyl, benzyl or C1-C3 alkylcarbonyl; X1, Y1, and Z1, are independently H, F, Cl, Br, I, C1-C4 alkyl, C1-C2 alkylthio or C1-C4 alkoxy; and	re
	55
p is 0, 1 or 2.  44. A method according to Claim 41 wherein the herbicide is a compound selected from the group	
consisting of	
	60

wherein 10

5

15

20

A is 
$$z_2$$
  $x_1$ 

ŧ

₫

25

*30* 

35

Rf is C1-C4 alkyl;

Rg is C1-C4 alkyl or C3-C6 cycloalkyl;

A1 is COORI, CH2OH or CHO;

R<sub>I</sub> is H; C<sub>1</sub>-C<sub>12</sub> alkyl optionally substituted by C<sub>1</sub>-C<sub>3</sub> alkyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl or phenyl; C<sub>3</sub>-C<sub>5</sub> alkenyl optionally substituted by phenyl or 1-2 C<sub>1</sub>-C<sub>3</sub> alkyl, F, Cl, Br or I; or C<sub>3</sub>-C<sub>5</sub> alkynyl optionally substituted by phenyl or 1-2  $C_1$ - $C_3$  alkyl, F, Cl, Br or I; B is H; C(O) $C_1$ - $C_6$  alkyl or C(O)phenyl optionally substituted by Cl, NO<sub>2</sub> or OCH<sub>3</sub>;

X2 is H, F, Cl, Br, I, OH or CH3;

Y2 and Z2 are independently H, C1-C8 alkyl, C1-C8 alkoxy, F, Cl, Br, I, Phenyl, NO2, CN, CF3 or SO2CH3;

X<sub>3</sub> is H, C<sub>1</sub>-C<sub>3</sub> alkyl, F, Cl, Br, I or NO<sub>2</sub>; and

L, M, Q and Rh are independently H, F, Cl, Br, I CH<sub>3</sub>, OCH<sub>3</sub>, NO<sub>2</sub>, CF<sub>3</sub>, CN, N(CH<sub>3</sub>)<sub>2</sub>, NH<sub>2</sub>, SCH<sub>3</sub> or SO<sub>2</sub>CH<sub>3</sub> provided that only one of M or Q may be a substituent other than H, F, Cl, Br, I, CH<sub>3</sub> or OCH<sub>3</sub>.


40


45

50

55

60








FIG.3

10 CTGCAGGTCGACTCTAGTGTACAAA	30 CATTGGGATGTGAAGGCTCAA	50 GGATGTGAATTGATACTCT	70 CATCAGGGGGGAGTTAATACG	90 IGHIGHACTCTHHITICCTHA	110 CAAGATTTTGACCCA
130 CTGGGTTTTTCTTGCAAGGTTTTT	150	170	190	210	270
250 CCAGCCACATTATETATGGACATC	270 CAAGGGGGAGTGTTATAAGAA	290 MAATCAAATTATGGTGGAT	310 GTCTACTCTTŒTŒATGAT	330 ETTETEAAATGETTAATGACA	350 TATTCAATGACATAT
370 TTCTATCCTTAATCACATATTTTTC	390 CITCACTITICATGCCTATAT	410 MAGGCCTTGTAATAGATA	430 GAAAAATACAAATAAATTEAAS	450 GAAGAAATAAAAATCTCTTAT	470 CTCTATATTTCTTAG
490 CITGITITITITITITITITATATIO	510 STACITTGAGCTATATTICA	530 TAACAGCATTCACATTCIT	550 ITICEATAGICTITEITCCCT	570 ITTTATATTTTAATTTACTGA	590 . AGTAACAAATACTTC
CACITICITICITICITICACCUTICA	630 TAMATATATCCAACATCTCA	650 ITTITICTITICCCCAATIC	670 TCAGACATTITTAATCTTTCTT	690 ITTETATTTATTTTCTTCATA	710 TTTIGATETETETE
CYTHICITICATICCATTTTTCCCTV	750 ITTCACGTGAATTCAATCAAG	770 TAGGACCETTTCAGTTTCE	790 TOGGGGCTCTCGTCTTCTCAGC	810 CTIANTATAMACEACCACA	830 CACCATCTACATTEC
850 CETTTCCTTTCAGTTTCCTCTCTC	870 ETECTCTCATTCAACAATAA M	890 TOGCOGCOGCTGCGGGGGCGCCGCCC	910 ICCATCTCCCTCTTTCTCCAA PSPSFSK	930 WACCETATOSTOCTOCTOCT T L 8 S 8 S S	950 CEANATOCTOCHOCC C S S T L
970 TOCTOCCTACATCCACCTTCCCTTT	990	1010	1030	1050	1070
L P R S T F P F					
1090 TYTCCACTACCCAAAAAGTTTCCGA	1110 GACCEAAAAAGCCGAAACTT	1130 TOSTTTCCCSTTTTSCCCC	1150 TEACEAACCCAGAAAGGGTTG	1170 COACETTCTOSTOGAGGCCC	1190 TEAAACAGAAGGG
S T T Q K V S E	TOKAETF	VERFAP	DEPREGS	DATAEVT	EREGY
1210 TTACCCACCTTTTTGCCTACCCACC					
TBVFATPG					
ACCETTACECACCCCACCCCCTT G Y A R A T G F	1350 CCCCCCCTTTCCATTCCCA P C V C I A T	1370 CCTOCCCCCCCCCCCACC S G P G A T	1590 PATCTCETCAETGGCCTCGC W L V S G L A	1410 EGACECCCTACTEGATAGCG1 DALLDSV	1430 ICCCCATIGITICETA PIVAI
1450 TAACCEGTCAAGTTGCACETAGGAT	1470 GATGGGTACTGATGCTTTTC	1490 . REGARACTECEATTETTERS	1510 ETAACTAGATCGATTACGAA	1530 GCATAATTATCTCGTTATGG	1550 COTAGAGGATATTC
T G Q V A R R M	15-70 T 5-70	1610	1630	1650	1670
R V V R E A F F	CETTECEAGATCGGGCCGGC	CTECCCCTGTTTTCATTCAT	GTACCTAAGGATATTCAGCA	ACAATTGGTGATACCTGACTG	CGATCAGCCAATGA
1690 GGTTGCCTGGTTACATGTCTAGGTT L P G Y N S R L					
1810	1830	1850	1870	1890	1910
S S E E L R R F					
1930 ATGGTACTGTTTATGCTAATTATGC	1950 TIGTGGACAGTAGTGATTTAT	TECTOSCATTICGGGTGAG	. 1990 STITGATGATAGAGTTACTGG	2010 AMETTAGMECTTTTECTA	2030 CCCCACCCAAAATTG
6 T V Y A N Y A	2070	2090	2110	2130	2150
TTCACATIGATATIGATICAGCIGA	GATTGGAAAGAACAAGCAGC	CICATETTICCATTIGICC	GGATATCAAGTTGGCGTTAC	LEGETTTEAATTEGATATTEG	AGAGTAAGGAAGGTA
2170	2190	2210	2230	2250	2270
L C L D F S A V	R Q E L T V Q	K V K T P L	EAATTTAAAACTTTTGGTGA # F K T F G D	A I P P Q Y A	I Q V L D
2290 ATGAGTTAACTAATGGGAGTGCTAT	2310	2330 AGCACCAGATETGGGCTGC	2350 CAATATTATAAGTACAGAA	2370	2390
ELTMGSAI					
2410 CCATCCCATTICCTTICCCCCCTCC N G F G L P A A					
2520 AGGTGGAGAATCTCCCAGTTAAGAT VEWLPVEI	2550 ITATGTTACTGAATAATCAAC M L L M H G M	2570 ACTICOCCAATGGICGGICA L G N V V O	2590 ATTOGRAGGATCOGTTCTATAR L E D R F Y K	2610 AGGETAACAGAGCACACACATA A M R A H T Y	2630 ACCTGGGGAATCCTT L G W P S
2650 CTANTCAGGCCGAGATETTTCCTAA W E A E I F P N					
2770 CTGGGCCATACTTGTTGGATGTGAT	2790 TGTACCTCATCAGGAACATG	2810 FICTACETATGATTCCCAG	2830 GGCCGGGCTTTCAAAGATGT	2850 GATCACAGAGGGTGACGGGA	2870 CAAGTTCCTATTGAC
C P Y L L D V 1	V P H Q E H V	L P M I P S	GGAFKDY	1 T E G D G R	5 5 Y .
TTIGAGGIGCTACAGAGCTAGTICE				FIG.4	

						-
10 GGATCCCTCT	TTCATTIGTTCTCATC	30 CATTITIGEGATICATGIGE	50 ATTTAATCAGTAGGACCCCS	70 TTTTAGCTTAGTAGTGCTCT	90 CATGTTCTCAACTTAATATT	110 MACCAACCACACT
130 CCATCTGCA1	TACCCTCCTTCCAGTT	150 TEGTETETECCTGCCCTCCC	170 ETTCAACAATGGCGGCGGCG	190 GCTCCATCTCCCTCTTCTTC A. P. S. P. S. S. S.	210 CGCTTTCTCCMAACCCTAT	230 מבכרון מכו ככו ככ
250	) September 1 AGANG	270		310 CCACCCCTCCACCTCACCCA	330	350
T S S 1	LLPRS	T F P F P H H	P K T T P	PPLHLTH	T W I H I H S	9 R R R
TICACCATAL	CEAATGTCATTTCCAC	TAACCAAAAGTTTCCCAGA	CCGAMMACCGMACTTTC	V S R F A P D	CGAACCEAGAAAGGGTTCCC E P R K G S D	VLVE
490 CCTCTCGAAA A L E 6	GAGAAGGGGTTACGGA	S10 CGTCTTTGCGTACCCAGGTG V F A Y P G E	530 GCGCTTCCATGGAGATTCAC A S M E I M	550 CAAGCTTTGACCCGTTCAAG Q A L T R S S	S70 CATCATCCGCAACGTGCTGC IIRKYLP	590 CACGTCACGAGCAG R N E Q
610 CCCCGTGTCT	TOCCOCCICACCCTTA	630	650 CCCGCCGTTTGCATTGCCACX	670 STETGGCCCCGGCGCGCCACCAA S G P G A T N	690 TCTCGTCAGCGGCCTCGCTG	710 ACCCCCTACTGGAT
77		750	770	790 GAAACTCETATTGTTGAGGT	<b>510</b>	830
S V P I	I V <sub>.</sub> A 1 T G	9 V Q R R M I	GTDAFQ	ETPIVEV	7 R S 1 T K E	W Y L Y
ATCCACCTA	CACCATATTECTAGGGT	TGTACGTGAAGCTTTTTTC	TOGOGAGATOGGGCCGGCC	COCCCTATTITGATTGATGT	PKDIQQQ	LVIP
971 GACTGGGAT	PACCEALTCACCTTACC	990 HEGITACATETETAGGITGS GYNSRLF	1010 ECTAMATICCCCAATGAGAT( C	1030 ECTTTTAGAACAAATTGTTAG L L E D I V R	CCTTATTTCTGAGTCAAAGA	1070 ACCCTGTTTTGTAT PVLY
109 GTGGGGGGT V G G	COCTOTTCCCAATCCAC	1110 TEAGGACTTGAGACGATTCC EDLRRF\	1130 TOGAGETEACGGÉTATCCC / E L T G 1 P	1150 CGTGGCAAGTACTTTGATGGG V A S T L M G	1170 TICTTGGAGETTTTGEAACTG LGAFPTG	1190 CCCATGAGCTTTCC DELS
121 CTTTCAATG L S H	TYCCCTATCCATCCTA	1230 CTGTTTATGCTAATTATGCTC V Y A N Y A N	1250 TI GGACASTAGTGATTIGTT / D S S D L L	1270 CCTCGCATTTGGGGTGAGGTT L A F G V R F	TGATGATAGAGTTACTGGAA	1310 ACTTAGAAGCTTTT LEAF
133 GCTAGCOGA A S R	CCAAAAATTGTTCACA	1350 TTGATATTGATTCAGCTGAG D I D S A E 1	1370 ATTGGAAAGAACAAGCAGCC   G K N K O P	1390 TCATGTTTCCATTTGTGCACA E V S 1 C A D	TATCAAGTTGGCGTTACAGG	1430 ETTTEAATTCGATA E L W S I
145 ETGGAGAGT L E S	ARCCARCETARACTOR	1470 AGTTGGATTTTTCTGCTTGG LDFSAU	1490 AGGCAGGAGTTGACGGAGCA L Q E L T E Q	1510 CAAAGTGAAGCACCCATTGA KVKEPLK	1530 ACTITAAAACTITIGGTGATG FKTFGDA	1550 CAATICCTCCGCAA - LIPP 0
157 TATGCTATO Y A 1	CACCTYPTACATCACT	1590 TAACTAATGGGAATGCTATTI T H G N A L	1610 ATAAGTACTGGTGTGGGGCA ISTGVGG	1630 ACACEAGATGTGGGCTGCTCI H Q M L/ A A Q	1650 NATACTATANGTACAGAAAGG YYKYRK	1670 TACGCCAATGGTTG
169 ACATCTGGT T S G	COLTRACCACCALTCC	1710 GATTIGGTTIGCCCGCTGCT F G L P A A	1730 Attegteceectettegaag I G A A V G R	1750 ACCEGATGAAGTTGTGGTTG PDEVVVD	1770 ACATTGATGGTGATGGCAGTT 1 D G D G S I	1790 TCATCATGAATGTG FINNV
			•			
16 CAGGAGCT Q E L		1830 CAGAATCTCCCAGTTAAGATT E N L P V K I	1850 PATGTTACTGAATAATCAAĆ K L' L W N D N	1870 ACTTGGGAATGGTGGTTCAAT L G N V V Q G	1890 IGGGAGGATCGGTTCTATAAG I E D R F Y K	1910 ECTANCAGAGCACAC A N R A N
19	30	1950	1970	1970 CTTGTGGCGTACCTGCTGCA C G V P A A I	2010 ACACTGACACATAGGGATGAT	2030 TCTTAGAGCTGCCATT L R A A I
	50 GTTAGACACTCCTGGG LDTPG	2070 CEATACTIGITGGATGTGAT PYLLOVI	2090 TETACCTCATCAGGAACATG V P N Q E N V	2110 TITYACCTATGATTCCCAGT	2130 GEOGRAGETTTCALAGATGT G G A F K D V	2150 CATCACAGAGGGTGAC 1 T E G D
CGGAGAAC	TO TO THE TOTAL TO	2190 CAAGCTACAGAGCTAGTTCT	2210 AGGCCTTGTATTATCTAAAA	2230 TAMACTYCTATTANGCCANA	2250 CATGUTCTGTCTATTAGTTT	2270 GTIGTTAGTTTTTGCT
GTGGCTT1	290 IGCTCGTTGTCACTGTT	2310' GTACTATTAAGTAGTTGATA	2330 TTTATGTTTGCTTTAAGTTT	2350 TGCATCATCTCCCTTTGGTT	2370 TTGAATGTGAAGGATTTCAG	2390 CAAAGITTCATTCTCT
24	.10	2430	2450	2470	2490 TTTCCTAACCTATGTTAGGC	2510 TGGGTCAGATTGGAAC

	208	
FIG. 6  119 122	194 197  ———————————————————————————————————	255 257
10  3.	A. GASHEHOLITISSI IRHVPRHEGGGYAEGYAATGFGVEIATGFGATHUSGLABALDSVPIVATGGVPRAHIGTOAFGETP IVEVTRSITKHWILWOVED PRWARAFF    IIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIII	250 350 350 350 350 350 350 350 350 350 3
4 8 0 0 8 6 6	а в с с и и с	4 B C C B F G





17.0   17.0	FIG. 6  381	586 59'5	
	410 GIVYANTANOSSOLLLAFGVRFOORVTGKLEAFASRAKIVNIO IOSAELGKIKOPHVSÍCLO KLALGCLUSILESKEGKLKLOFSARGELVOKKVPLHFKÍF GIVYANTANOSSOLLLAFGVRFOORVTGKLEAFASRAKIVNIO IOSAELGKIKOPHVSÍCLO KLALGCLUSILESKEGKLKLOFSARGELFORKKPLHFKÍF GIVYANTAVOSSOLLLAFGVRFOORVTGKLEAFASRAKIVNIO IOSAELGKIKOPHVSÍCLO KLALGCALNILESKEGKLKLOFSARGELFORKKPLHFKÍF GIVYANTAVEHSOLLLAFGVRFOORVTGKLEAFASRAKIVNIO IOSAELGKKTPHVSVCGOVKLALGGGWRKTLAFGGKRRGELFORKKPLHFFFF GIVTANTAVEHSOLLLAFGVRFOORVTGKLEAFASRAKIVNIO IOSAELGCHVALGGGWRKTLAGGGARKTLAGGARKTLAGGARKTLAGGARKTLAGGARKTLAGGARKTLAGGARKTLAGGARTTHLGGGARKTLAGGARKTLAGGARTTHLGGGARTTHLGGGARKTLAGGARTTHLGGGARTTHLAGGARTTHLGGGARTTHLGGGARTTHLGGGARTTHLGGGARTTHLGGGARTTHLGGGARTTHLAGGARTTHAGGARTHAGAAGGA		######################################



È

Ī





. \$

i

Publication number:

**0 257 993** A3

**②** 

## EUROPEAN PATENT APPLICATION

2) Application number: 87307384.5

② Date of filing: 20.08.87

(5) Int. Cl.5: C07H 21/04 , C12N 15/00 , C12N 9/88 , C12N 1/20 , A01H 1/00 , A01H 5/10 , A01N 25/00 , A01N 47/36

Priority: 26.08.86 US 900609

Date of publication of application:02.03.88 Bulletin 88/09

Designated Contracting States:

AT BE CH DE ES FR GB GR IT LI LU NL SE

Date of deferred publication of the search report: 02.05.90 Bulletin 90/18

Applicant: E.I. DU PONT DE NEMOURS AND COMPANY
 1007 Market Street
 Wilmington Delaware 19898(US)

Inventor: Bedbrook, John Robert 110 Cambridge Way Piedmont, California 94611(US)
Inventor: Chaleff, Roy Scott

1 Spring Lane
Chadds Ford, Pennsylvania 19317(US)
Inventor: Falco, Saverio Cari
1902 Millers road
Arden, Delaware 19810(US)
Inventor: Mazur, Barbara Jean
4601 Weldin road
Wilmington, Delaware 19803(US)
Inventor: Yadav, Narendra Singh
127 Jade Drive
Wilmington, Delaware 19810(US)

(2) Representative: Hildyard, Edward Martin et al Frank B. Dehn & Co. Imperial House 15-19 Kingsway London WC2B 6UZ(GB)

(S) Nucleic acid fragment encoding herbicide resistant plant acetolactate synthase.

(F) A nucleic acid fragment encoding a herbicideresistant plant acetolactate synthase protein is disclosed. This nucleic acid fragment contains at least one nucleotide mutation resulting in one amino acid change in one of seven substantially conserved regions of acetolactate synthase amino acid homology. This mutation results in the production of an acetolactate synthase protein which is resistant to sulfonylurea herbicide compounds compared to the wild-type protein. Transformation of herbicide sensitive plants or plant cells with the fragment results in resistance to the herbicide.

	10 30	30	70	90	110
A.	MALAAAAPEPEFEETLESSESSTLUPTST	PEPEEPECT POPEELTPI KI		IGEAR LIARES VACENERS	SOULVEALERECVIOVIATES
	11014 6126 111118 [11 131116[1]	12206211111211111   13	111811111111111111111111111111111111111		SOAFAE WEEKALDALVEL
В.	MOTOR* 1635-835-W. 321 C3. 2541 ca. ce. ce.	PPPERPERIT PPPERINTHENS	REGESTALL CENAL PLANTARY	1 111 111111 111	i de de la
	(11) (1) (1) (1) (1)	!! !!		PRESCRIPTION OF THE PROPERTY O	ADILVEALEROGYETYPATPO
c.		DESCRIPTIONS SERVICE (CESSO)		1	
_	MIROSILKHFAIKRCFORIATIRIPHRESV		ACRES COMPANDED	APPLRATEPONOTSFVGLT	COLL NEWS ENDING IN GIRG
D.	WINGZ LIKELY I KECLOFI YANE PAYA	Christ a system i mentioner			
-				MÁSSOTTSTRIALFÍ	BAEI IVHILEOGSIKIVIĞIPĞ
E.					H H H H <u>J C H</u>
_				KCB	CACUVATAL BACCVISTYF CTPG
P.					11 (1 ) 111 (1111)

119 122 — E<sub>2</sub> E<sub>1</sub> Sub-sequence A

FIG.6

Xerox Copy Centre



## EUROPEAN SEARCH REPORT

EP 87 30 7384

<del>-</del>		DERED TO BE RELEVAN	Relevant	CLASSIFICATION OF THE
Category	Citation of document with in- of relevant pas		to claim	APPLICATION (Int. CL4)
х о,х	SCIENCE, vol. 224, 2 pages 1443-1445; R.S "Acetolactate syntha action of two sulfor in higher plants" * Whole document * ULCA SYMP. SER., vol MOLECULAR STRATEGIES	9th June 1984, i. CHALEFF et al.: ise is the site of iylurea herbicides  48, UCLA SYMP. is FOR CROP th March - 6th April is Steamboat Springs et al.: arieties resistant	30	C 07 H 21/04 C 12 N 15/00 C 12 N 9/88 C 12 N 1/20 A 01 H 1/00 A 01 H 5/10 A 01 N 25/00 A 01 N 47/36
0 <b>,</b> A	IDEM		1-29,31 -44	·
0,Y	WORLD BIOTECH REPORT PROCEEDINGS OF BIOTE DC, October 1985, pa MAZUR et al.: "Clon- resistance genes in- plants" * Whole document *	ECH'85, Washington, ages 97-108; B.J. Ing herbicide	1-44	TECHNICAL FIELDS SEARCHED (Int. CL4)  C 07 H C 12 N A 01 H A 01 N
0,Y	TRENDS IN GENETICS, 89-93; M.A. ESTELLE mutants of Arabidop * Page 91, paragrap	et al.: "The sis"	1-44	
	The present search report has b	een drawn up for all claims		
	Place of search	Date of completion of the search	MARK	Examiner
TH	E HAGUE	12-01-1990	MADI	OOX A.D.
Y: pa do A: te	CATEGORY OF CITED DOCUMENTS  X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: Intermediate document CATEGORY OF CITED DOCUMENTS  T: theory or principle underly after the filing date D: document cited in the app L: document cited for other CATEGORY OF CITED DOCUMENTS  E: earlier patent document after the filing date D: document cited in the app L: document cited for other CATEGORY OF CITED DOCUMENTS  E: earlier patent document			lished on, or

Office

Page

EP 87 30 7384

<u> </u>		DERED TO BE RELEVA			
Category	Citation of document with in of relevant pas	dication, where appropriate, ssages		levant claim	CLASSIFICATION OF THE APPLICATION (Lot. Cl. 4)
Υ	NUCLEIC ACIDS RESEAU 11, 1985, page 4011 "Nucleotide sequence gene which encodes a synthase" * Whole document *	S.C. FALCO et al.: e of the yeast ILV2	1-4	4	
P,X	NATO ASI SER. A, vo BIOL.) edited by Di- and Nam-Hai Chua, P ADVANCED STUDY INST MOLECULAR BIOLOGY, June 1987; B.J. MAZI "Acetolactate synthe enzyme of the sulfo * Whole document *	ter von Wettstein ROCEEDINGS OF NATO ITUTE ON PLANT Cansberg, 10th-19th UR et al.: ase, the target	1-4		
0,A	BIOLOGICAL ABSTRACT C.R. SOMERVILLE et herbicide resistanc level" * Whole abstract *	S/RMM, BR30:49417; al.: "Selection for e at the whole plant	1-4	4	TECHNICAL FIELDS SEARCHED (Int. CL4)
A	EP-A-0 154 204 (MD * Whole document *	LECULAR GENETICS)	1-4	14	
A	PROC. NATL. ACAD. S June 1986, pages 44 al.: "Single amino in the enzyme aceto confer resistance t sulfometuron methyl * Abstract *	18-4422; N. YADAV et acid substitutions lactate synthase o the herbicide	1-4	14	
E	EP-A-0 292 435 (CI * Pages 37,38 *	BA-GEIGY)		29-31 3,34,	
	The present search report has t	een drawn up for all claims			
TH	Place of search E HAGUE	Date of completion of the search 12-01-1990		MADI	Examiner DOX A.D.
THE HAGUE  CATEGORY OF CITED DOCUMENTS  X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document  T: theory or principle underlysh E: earlier patent document, bu after the filing date D: document cited in the applil L: document cited for other residuence document  &: member of the same patent document			t, but publ application er reasons	ished on, or	